

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/CS/29 Forms of Magnesium - S. Wheat

Rothamsted Research

Rothamsted Research (1971) *70/W/CS/29 Forms of Magnesium - S. Wheat* ; Numerical Results Of The Field Experiments 1970, pp 176 - 177 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

FORMS OF MAGNESIUM

(70/W/CS/29)

Effect of forms of magnesium fertiliser on spring wheat - Woburn
Stackyard C, 3rd year.

For details of treatments etc. and previous years' results see
'Results' 68/C/35 and 69/W/CS/29. Mg and limestone treatments
are cumulative.

Area of each plot: 0.0072. Area harvested: 0.0032.

Basal applications: 540 lb (20:10:10). Weedkiller: Ioxynil at
7.5 oz and mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.:- Ploughed: 12 Nov, 1969. NPK applied:
26 Mar, 1970. Mg and limestone treatments applied: 31 Mar.
Power harrowed, seed drilled at 175 lb: 1 Apr. Weedkiller
applied: 18 May. Combine harvested: 14 Aug. Variety:
Kolibri.

Standard error per plot.

Grain, cwt: 3.21 or 31.6% (21 d.f.)

70/W/CS/29

SUMMARY OF RESULTS

O	D2	E	C1	C2	DL	L2	EL	Mean
GRAIN: CWT								
(±1.61)								
11.0	9.9	11.3	10.0	9.4	11.2	6.9	11.6	10.2
STRAW: CWT								
6.6	6.5	7.7	6.2	6.6	6.8	5.0	6.2	6.4

Mean D.M. %: Grain: 82.0
Straw: 90.1