

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/CS/45 Nematicides in Rows - Barley

Rothamsted Research

Rothamsted Research (1971) *70/W/CS/45 Nematicides in Rows - Barley* ; Numerical Results Of The Field Experiments 1970, pp 191 - 192 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

NEMATICIDES IN ROWS

(70/W/CS/45)

Residual effects of nematicides in rows and irrigation, applied to sugar beet 1969 - Woburn Butt Close II 1970, barley second year.
For previous year's results see 'Results' 69/W/CS/45.

Design: 3 blocks of 2 plots split into 8.

Area of each sub plot: 0.0069. Area harvested: 0.0069.

Treatments: All combinations of:-

1. Whole plots: Residue of irrigation 1969: None (0), full (C).
2. Sub plots: Residues of nematicides 1969: None (D0), 56 (D2), 84 (D3), 112 lb (D4) 'D-D' placed in rows before drilling. 56 lb (T) 'Telone' placed in rows before drilling, 1.5 (K1), 3.0 (K2), 4.0 lb (K3) aldicarb placed in rows at drilling.

NOTE: The nitrogen test to beans in 1968 is now ignored.

Basal applications: 400 lb (20:10:10). Weedkiller: Ioxynil at 7.5 oz plus mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.: Ploughed: 29 Dec, 1969. Power harrowed:

26 Mar, 1970. Seed combine drilled at 140 lb: 27 Mar.

Weedkiller applied: 16 May. Combine harvested: 25 Aug.

Variety: Julia.

Standard error per sub plot.

Grain, cwt: 2.18 or 8.1% (28 d.f.)

70/W/CS/45

SUMMARY OF RESULTS

GRAIN: CWT

	D0	D2	D3	D4	T	K1	K2	K4	Mean
(±1.26)*									
O	27.4	29.2	24.2	27.5	29.7	25.9	28.6	25.7	27.3
C	25.5	26.6	26.3	24.5	26.9	27.7	24.4	29.4	26.4
Mean (±0.89)	26.4	27.9	25.3	26.0	28.3	26.8	26.5	27.6	26.9

Mean D.M. %: 83.6

* For use in horizontal and interaction comparisons only