

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



**ROTHAMSTED
RESEARCH**

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/WW/5 Winter Wheat Cultivations and Bulb Fly

Rothamsted Research

Rothamsted Research (1971) *70/R/WW/5 Winter Wheat Cultivations and Bulb Fly* ; Numerical Results Of The Field Experiments 1970, pp 221 - 222 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-59>

WINTER WHEAT

(70/R/WW/5)

Cultivations and bulb fly (*Leptohylemyia coarctata*) -
Stackyard 1970.

Design: 4 randomised blocks of 3 plots.

Area of each plot: 0.0362. Area harvested: 0.0129.

Treatments: Tine cultivation, no ploughing (C), normal ploughing
(P), scraped and ploughed (SP).

NOTE: The SP treatment was done by a special plough, the top
2-3 inches of soil being scraped into the open furrow
just ahead of the plough.

Basal applications: 280 lb (8:20:16) combine drilled, 0.84 cwt N
as 'Nitro-Chalk' top dressed in spring. Weedkiller: Ioxynil
at 9 oz and mecoprop at 27 oz in 20 gals.

Cultivations, etc.: Subsoiled 28 inches apart, 18 inches deep:
4 Aug, 1969. C plots deep-tine cultivated, P plots
ploughed, SP plots scraped and ploughed: 11 Sept. All
plots rotary cultivated: 11 Nov. Seed combine drilled at
180 lb: 14 Nov. 'Nitro-Chalk' applied: 28 Apr, 1970.
Weedkiller applied: 6 May. Combine harvested: 27 Aug.
Variety: Cappelle. Previous crops: Winter wheat 1968,
Fallow 1969.

NOTE: Samples were taken for bulb fly eggs in February 1970.

Standard error per plot.

Grain, cwt: 3.15 or 10.6% (6 d.f.)

70/R/VV/5

SUMMARY OF RESULTS

GRAIN: CWT

C	P	SP	Mean
	(±1.58)		
24.9	31.8	32.4	29.7

Mean D.M. %: 83.6