

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/WW/7 Winter Wheat Seed Dressings and Soil-borne Diseases

Rothamsted Research

Rothamsted Research (1971) *70/R/WW/7 Winter Wheat Seed Dressings and Soil-borne Diseases ; Numerical Results Of The Field Experiments 1970*, pp 225 - 226 - DOI:
<https://doi.org/10.23637/ERADOC-1-59>

WINTER WHEAT

(70/R/WW/7)

Seed dressings and soil-borne diseases, Highfield Drive 1970.

Design: 4 randomised blocks of 8 plots.

Area of each plot: 0.0050. Area harvested: 0.0027.

Treatments: None (0) (2 plots per block) and all combinations of:-

1. Seed dressings (as wettable powders): 50% benomyl (B), 75% 'Plantvax' (P), 60% thiabendazole (T).
2. Rates of application per 63 lb of seed: 1 oz (1), 4 oz (2) of wettable powder.

Basal applications: 280 lb (8:20:16) combine drilled, 1.0 cwt N as 'Nitro-Chalk' top dressed in spring. Weedkillers: Paraquat at 0.75 lb ion in 20 gals, 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc.: Paraquat applied: 17 Sept, 1969. Ploughed: 22 Sept. Rotary cultivated: 15 Oct. Seed combine drilled at 160 lb: 27 Oct. 'Nitro-Chalk' applied: 28 Apr, 1970. 2,4-D/dichlorprop applied: 8 May. Combine harvested: 27 Aug. Variety: Cappelle. Previous crops: Barley 1968, winter wheat 1969.

NOTE: Samples were taken for root and foot rot diseases.

Standard error per plot.

Grain, cwt: 2.47 or 6.6% (21 d.f.)

70/R/W/7

SUMMARY OF RESULTS

GRAIN: CWT

	O	B	P	T	Mean
			(± 1.24)		(± 0.71)
1		37.6	38.0	39.2	38.3
2		37.4	37.9	37.7	37.7
Mean (± 0.87)	36.0	37.5	38.0	38.5	37.5*

* General mean

Mean D.M. %: 84.3