

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Numerical Results of the Field Experiments 1970

[Full Table of Content](#)

International Agricultural Research  
Institute  
London Agricultural Station  
Woburn, Bedfordshire  
MK42 9EF  
ENGLAND  
01235 533111  
www.rii.ac.uk  
© 2012 Rothamsted Research. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without the prior written permission of Rothamsted Research. For more information, please contact the Rothamsted Research Library, Rothamsted, Harpenden, Herts, AL5 2JQ, UK. Tel: +44 (0)1494 473900. Fax: +44 (0)1494 473901. Email: [library@rothamsted.ac.uk](mailto:library@rothamsted.ac.uk)

## 70/R/WS/3 Spring Wheat Ccc (Ccc) in Grain

### Rothamsted Research

Rothamsted Research (1971) *70/R/WS/3 Spring Wheat Ccc (Ccc) in Grain* ; Numerical Results Of The Field Experiments 1970, pp 235 - 236 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

SPRING WHEAT

(70/R/WS/3)

CCC\* in grain, Long Hoos V 1970.

\* Chlormequat.

Design: 4 randomised blocks of 4 plots, split into 3.

Area of each sub-plot: 0.0096. Area harvested: 0.0064.

Treatments: All combinations of:-

Whole plots: 1. CCC in grain: None (G0), sprayed the previous year with 2 lb chlormequat in 40 gals just after ear emergence (GG).

2. CCC spray in spring 1970: None (S0), 1.5 lb chlormequat in 35 gals on 28 May, 1970 (SS).

Sub-plots: 3. Nitrogen: 0.6 (N1), 1.2 (N2), 1.8 (N3) cwt N as 'Nitro-Chalk'.

Basal applications: 225 lb (0:20:20) combine drilled. Weedkiller: 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc.: Deep-tine cultivated: 27 Oct, 1969. Seed combine drilled at 170 lb: 19 Apr, 1970. N applied: 28 Apr. Weedkiller applied: 27 May. Combine harvested: 28 Aug. Variety: Kolibri. Previous crops: Spring oilseed rape 1968, potatoes 1969.

NOTE: Crop samples were taken from 2 blocks on 14 Aug for measurements of height and components of yield.

Standard errors per plot. Grain, cwt:

Whole plot: 1.46 or 4.9% (9 d.f.)

Sub plot: 2.57 or 8.7% (24 d.f.)

70/R/WS/3  
SUMMARY OF RESULTS

GRAIN: CWT

	SO	SS	N1	N2	N3	Mean
	(±0.73)		(1) and (2)			(±0.51)
GO	29.8	29.5	29.3	29.0	30.8	29.7
GG	29.4	30.0	27.7	30.2	31.1	29.7
			(1) and (2)			(±0.51)
		SO	29.1	29.3	30.5	29.6
		SS	27.9	29.9	31.4	29.7
Mean (±0.64)			28.5	29.6	30.9	29.7

Mean D.M. %: 85.9

(1) (±0.90) For use in vertical and diagonal comparisons only

(2) (±0.91) For use in horizontal and interaction comparisons only

STRAW: CWT

	SO	SS	N1	N2	N3	Mean
GO	13.8	11.7	11.3	12.9	14.0	12.8
GG	12.7	11.8	12.1	12.1	12.5	12.2
		SO	12.8	12.7	14.1	13.2
		SS	10.6	12.3	12.3	11.8
Mean			11.7	12.5	13.2	12.5

Mean D.M. %: 82.6