

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/B/2 and 70/W/B/2 Barley Varieties X N and Mildew Control (Erysiphe Graminis)

Rothamsted Research

Rothamsted Research (1971) *70/R/B/2 and 70/W/B/2 Barley Varieties X N and Mildew Control (Erysiphe Graminis)* ; Numerical Results Of The Field Experiments 1970, pp 247 - 254 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

BARLEY

(70/R/B/2 and 70/W/B/2)

Varieties x N and mildew control (*Erysiphe graminis*), Rothamsted (R)
Great Knott I and Woburn (W) Great Hill III 1970.

Design: Great Knott I (R): 4 randomised blocks of 14 plots, split
into 3.
Great Hill III (W): 4 randomised blocks of 16 plots, split
into 3.

Area of each sub plot: Great Knott I (R): 0.0080. Area harvested: 0.0054.
Great Hill III (W): 0.0048. Area harvested: 0.0032.

Treatments: All combinations of:-

Whole plots: 1. Varieties: Akka (A), Bala - Great Hill III (W) only
(B), Julia (J), Midas (M), Sultan (S), Vada (V),
Zephyr (Z), Akka sown in late May (AL). Seed rates:
Great Knott I (R) - A and AL plots 150 lb, remainder
140 lb. Great Hill III (W): All plots 140 lb.

2. Fungicide: None (O), 1 lb ethirimol as a seed
dressing (F).

Sub plots: 3. Nitrogen: 0.3 (N1), 0.6 (N2), 0.9 (N3) cwt N (R)
and 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N (W) as
'Nitro-Chalk'.

Basal applications: 220 lb (0:20:20) combine drilled. Weedkillers:-
Great Knott I (R): 2,4-D at 0.5 lb and dichlorprop at 2 lb in
20 gals. Great Hill III (W): Diquat at 0.34 lb and paraquat at
0.34 lb in 25 gals. Ioxynil at 9 oz and mecoprop at 27 oz in
25 gals.

Cultivations, etc.:-

Great Knott I (R): Ploughed: 17 - 31 Oct, 1969 and 3 Nov. Combine
drilled (except AL plots): 2 Apr, 1970. N applied: 14 Apr.
Weedkiller applied: 18 May. AL plots: N applied, seed combine
drilled: 19 May. A plots combine harvested: 30 July. Remaining
plots* combine harvested: 14 Aug. Previous crops: Fallow 1968,
winter wheat 1969.

Great Hill III (W): Ploughed: 4 Sept, 1969. Rotary cultivated:
24 Oct. Diquat/paraquat applied: 17 Mar, 1970. Combine drilled
(except AL plots): 27 Mar. All N applied: 9 Apr. Ioxynil/mecoprop
applied: 15 May. AL plots combine drilled: 20 May. A plots
combine harvested: 31 July. Remaining plots excluding AL
combine harvested: 12 Aug. Previous crops: Potatoes 1968, fallow
1969.

70/R&W/B/2

- NOTES: (1) The weather after sowing AL plots was very dry, growth was very poor and yields were not taken.
(2) Samples were taken for assessment of mildew (*Erysiphe graminis*) and other foliar diseases.
(3) At Woburn the appearance of the crop in July caused doubt about the allocation of the N-rates to sub-plots within some of the whole plots of block 1: this block has been omitted from the tables involving N and from the corresponding analysis.

Standard errors per plot.	Grain, cwt:
Great Knott I (R):	Whole plot: 1.58 or 5.4% (33 d.f.) Sub plot: 2.42 or 8.3% (72 d.f.)
Great Hill III (W)	
All 4 blocks:	Whole plot: 4.93 or 14.0% (39 d.f.)
Block 1 omitted:	Whole plot: 4.69 or 14.6% (26 d.f.) Sub plot: 5.08 or 15.9% (56 d.f.)

70/R&W/B/2

SUMMARY OF RESULTS

GRAIN: CWT

GREAT KNOTT I (R)

	A	J	M	S	V	Z	Mean
	(±0.79)						(±0.32)
O	14.9	33.3	32.0	31.5	30.1	32.1	29.0
F	15.4	33.2	31.6	32.9	30.7	33.6	29.6
	(1) and (2)						(±0.35)
N1	13.8	29.3	27.2	28.6	27.8	28.4	25.9
N2	14.5	34.2	33.7	33.7	30.5	33.7	30.1
N3	17.0	36.3	34.3	34.2	32.9	36.6	31.9
Mean (±0.56)	15.1	33.3	31.8	32.2	30.4	32.9	29.3
	O	F					
	(3) and (4)						
N1	25.2	26.5					
N2	29.5	30.6					
N3	32.2	31.6					

Mean D.M. %: 85.2

(1) (±0.89) (3) (±0.52) For use in horizontal and diagonal comparisons only

(2) (±0.86) (4) (±0.49) For use in vertical and interaction comparisons only

70/R&W/B/2

GRAIN: CWT

GREAT HILL III (W)

ALL 4 BLOCKS

	A	B	J	M	S	V	Z	Mean
	(±2.46)							(±0.93)
O	28.1	34.3	34.8	38.9	32.6	31.8	36.2	33.8
F	27.0	39.0	37.5	37.2	36.4	39.4	38.8	36.5
Mean (±1.74)	27.5	36.7	36.2	38.0	34.5	35.6	37.5	35.1

BLOCK 1 OMITTED

	(1) and (2)							(±0.78)
N1	21.1	32.1	27.0	34.2	30.0	31.3	28.2	29.1
N2	26.3	36.2	36.8	33.3	30.8	33.6	35.5	33.2
N3	25.6	30.0	31.9	35.4	38.3	37.4	38.0	33.8
Mean (±1.91)	24.3	32.8	31.9	34.3	33.0	34.1	33.9	32.0

	O	F
	(3) and (4)	

N1	28.4	29.8
N2	31.7	34.7
N3	31.9	35.7

Mean (±1.02)	30.7	33.4
--------------	------	------

Mean D.M. %: 82.6

(1) (±2.56) (3) (±1.37) For use in horizontal and diagonal comparisons only

(2) (±2.07) (4) (±1.11) For use in vertical and interaction comparisons only

BARLEY

(70/R/B/3 and 70/W/B/3)

Deep drilled urea and 'Nitro-Chalk', Rothamsted (R) Great Knott I and Woburn (W) Great Hill III 1970.

Design: 4 blocks of 8 plots (randomisation restricted) plus 1 nil plot per block.

Area of each plot:

Great Knott I (R): 0.0161.	Area harvested: 0.0103.
Great Hill III (W): 0.0154.	Area harvested: 0.0099.

Treatments: No nitrogen (NO) and all combinations of:-

1. Forms of nitrogen: 'Nitro-Chalk' 21% N (C), urea 46% N (U).
2. Methods of application: Deeply drilled by 'Fiona' drill in 6 inch rows before drilling seed (D), broadcast by hand after drilling seed (B).
3. Rates of nitrogen: 0.5 (N1), 1.0 (N2) cwt N.

NOTE: (1) The seed was drilled on all plots by the 'Fiona' drill in rows 6 inches apart.

Basal applications: Manures: 280 lb (0:20:20). Broadcast by 'Fiona' drill.

Weedkillers: Great Knott I (R): 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Great Hill III (W): 0.34 lb diquat plus 0.34 lb paraquat in 25 gals, ioxynil at 7.5 oz plus mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.:

Great Knott I (R): Ploughed: 17 - 31 Oct and 3 Nov, 1969. Seed drilled at 120 lb, basal PK, urea and 'Nitro-Chalk' applied: 2 Apr, 1970. Weedkiller applied: 18 May. Combine harvested: 12 Aug. Variety: Julia. Previous crops: Fallow 1968, winter wheat 1969.

Great Hill III (W): Ploughed: 4 - 10 Sept, 1969. Rotary cultivated: 24 Oct. Deep-tine cultivated: 1 Nov. Diquat/paraquat applied: 17 Mar, 1970. Seed drilled at 120 lb, basal PK, urea and 'Nitro-Chalk' applied: 31 Mar. Ioxynil/mecoprop applied: 15 May. Combine harvested: 11 Aug. Variety: Julia. Previous crops: Potatoes 1968, fallow 1969.

70/R&W/B/3

- NOTES: (2) The 'Fiona' drill was used on all plots to prepare the seedbed, being drawn idle through the E and NO plots.
(3) The percentage of N in the grain was determined.

Standard errors per plot. Grain, cwt:

Great Knott I (R): 1.23 or 3.5% (21 d.f.)
Great Hill III (W): 3.81 or 9.8% (21 d.f.)

70/R&W/B/3

SUMMARY OF RESULTS

GRAIN: CWT

GREAT KNOTT I (R)

	D	B	N1	N2	Mean
	(±0.43)		(±0.43)		(±0.31)
C	35.7	35.1	34.3	36.5	35.4
U	34.8	34.7	33.8	35.8	34.8
			(±0.43)		(±0.31)
		D	34.2	36.4	35.3
		B	33.9	35.9	34.9
Mean (±0.31)			34.0	36.1	35.1

NO 18.6
 General mean: 33.3

Mean D.M. %: 85.2

70/B&W/B/3

GRAIN: CWT

GREAT HILL III (W)

	D	B	N1	N2	Mean
	(±1.35)		(±1.35)		(±0.95)
C	39.4	40.1	40.7	38.8	39.8
U	38.8	37.8	38.9	37.7	38.3
			(±1.35)		(±0.95)
		D	40.0	38.2	39.1
		B	39.6	38.3	39.0
Mean (±0.95)			39.8	38.2	39.0

NO 29.5
 General mean: 38.0

Mean D.M. %: 80.2