Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1970



Full Table of Content

70/R/B/4 Barley Early and Late Mildew (Erysiphe Graminis)

Rothamsted Research

Rothamsted Research (1971) 70/R/B/4 Barley Early and Late Mildew (Erysiphe Graminis); Numerical Results Of The Field Experiments 1970, pp 255 - 256 - DOI: https://doi.org/10.23637/ERADOC-1-59

BARLEY

(70/R/B/4)

Early and late mildew (Erysiphe graminis), Long Hoos V 1970.

Design: 4 randomised blocks of 4 plots.

Area of each plot: 0.0257. Area harvested: 0.0129.

Treatments: Fungicide, ethirimol:
None

0.25 lb as seed dressing

2.0 lb as seed dressing plus 1 lb in 35 gals

as a foliar spray on 2 occasions

1.0 lb in 35 gals as a foliar spray on 2 occasions

(8D&S)

Basal applications: 360 lb (20:10:10) combine drilled. Weedkiller: 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc.: Deep-tine cultivated: 27 Oct, 1969. Seed combine drilled at 140 lb: 19 Apr, 1970. Weedkiller applied: 27 May. Fungicide sprays applied: 16 June and 8 July. Combine harvested: 12 Aug. Variety: Zephyr. Previous crops: Spring oilseed rape 1968, potatoes 1969.

NOTE: Samples were taken for assessment of mildew (Erysiphe graminis) and other foliar diseases and for assessment of tiller number, tiller length and ear sizes.

Standard error per plot.
Grain, cwt: 2.33 or 7.8% (9 d.f.)

SUMMARY OF RESULTS

0		ID	8D8s	8s	Mean
		PSI	GRAIN: CWT	A JOSEPH I	
(±1.17)					
28.5		29.5	31.1	29.8	29.7
			STRAW: CWT		in a es
15.0		15.5	17.0	15.3	15.7

Mean D.M. %: Grain: 82.0 Straw: 82.8