

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/S/B/2 Barley N Rates to Barley After Grass

Rothamsted Research

Rothamsted Research (1971) *70/S/B/2 Barley N Rates to Barley After Grass* ; Numerical Results Of The Field Experiments 1970, pp 264 - 265 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

BARLEY

(70/S/E/2)

N rates to barley after grass, Saxmundham Grove Plot 1970.

Design: 2 randomised blocks of 8 plots.

Area of each plot: 0.0046. Area harvested: 0.0018.

Treatments:- Nitrogen: None (N0), 0.4 (N1), 0.8 (N2), 1.0 (N3),
1.2 (N4), 1.4 (N5), 1.6 (N6), 2.0 (N7) cwt N as 'Nitro-Chalk'.

Basal applications: 560 lb (0:20:20) broadcast. Weedkiller:
Mecoprop at 36 oz and 2,4-D at 9 oz in 20 gals.

Cultivations, etc.: Ploughed: 30 Oct, 1969. N and PK applied,
seed drilled: 23 Apr, 1970. Weedkiller applied: 28 May. Combine
harvested: 18 Aug. Variety: Julia. Previous crops: 3 years
grass.

Standard error per plot.

Grain, cwt: 1.47 or 5.5% (7 d.f.)

70/S/B/2

SUMMARY OF RESULTS

NO	N1	N2	N3	N4	N5	N6	N7	Mean
GRAIN: CWT								
(±1.04)								
14.9	23.3	25.8	28.6	30.8	27.9	31.0	33.5	27.0
STRAW: CWT								
15.5	21.8	25.9	30.0	30.2	32.2	35.1	38.3	28.6

Mean D.M. %: Grain: 86.0
Straw: 88.4