

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/BE/7 Spring Beans Chemical Control of Soil-borne Pathogens

Rothamsted Research

Rothamsted Research (1971) *70/R/BE/7 Spring Beans Chemical Control of Soil-borne Pathogens ; Numerical Results Of The Field Experiments 1970*, pp 278 - 279 - DOI:
<https://doi.org/10.23637/ERADOC-1-59>

SPRING BEANS

(70/R/BE/7)

Chemical control of soil-borne pathogens, Barnfield (Valley Section, part of plot 3 and 4) 1970.

Design: 2 blocks of 2 plots split into 5.

Area of each sub plot: 0.0014. Area harvested: 0.0006.

Treatments: All combinations of:-

Whole plots: 1. Manuring: Classical treatments since 1968:-

Plot 3: no manures (O)

Plot 4: P, K, Na, Mg (30 lb P as superphosphate, 200 lb K as sulphate of potash, 80 lb Na as agricultural salt, 20 lb Mg as sulphate of magnesia) (PKNaMg)

Sub plots: 2. Chemical control:-

None (O)

Nematicide, 10% aldicarb ('Temik') at 10 lb a.i. (N)

Fungicide, 50% benomyl at 20 lb a.i. (F)

Insecticide, 50% gamma BHC at 2 lb a.i. (I)

Nematicide, fungicide and insecticide (as above) together. (NFI)

All plots were rotary cultivated immediately after application of chemicals.

Basal applications: Manures - none. Weedkiller: Simazine at 1 lb in 20 gals. Insecticide: Demeton-s-methyl at 3.5 oz in 40 gals.

Cultivations, etc.: P, K, Na, Mg applied: 21 Nov, 1969. Ploughed: 1 - 19 Dec. Chemical control treatments applied, all plots rotary cultivated: 20 Mar, 1970. Seed drilled at 200 lb: 20 Apr. Weedkiller applied: 24 Apr. Insecticide applied: 17 June. Harvested by hand: 3 Sept. Variety: Maris Bead.

NOTE: Wilted plants were counted during the season. Crop samples were taken for root disease assessment, and soil samples for counting *Sitona* beetle larvae. Counts were made of stem eelworm (*Ditylenchus dipsaci*) incidence.

Standard error per sub plot.

Grain, cwt: 2.44 or 15.1% (8 d.f.)

70/R/BE/7

SUMMARY OF RESULTS

GRAIN: CWT

	O	N	F	I	NFI	Mean
			(±1.72)*			
O	9.5	19.5	15.7	13.3	16.3	14.9
PK Na Mg	15.3	20.2	15.5	13.1	23.0	17.4
Mean (±1.22)	12.4	19.8	15.6	13.2	19.7	16.2

Mean D.M. %: 79.4

* For use in horizontal and interaction comparisons only