

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Numerical Results of the Field Experiments 1970

[Full Table of Content](#)

International Agricultural Research  
Institute  
London Agricultural Station  
Woburn, Bedfordshire  
MK42 9EF  
ENGLAND  
025  
© 1971 Rothamsted Research. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without the prior written permission of Rothamsted Research. Printed in Great Britain by the University Press, Cambridge.

## 70/R/P/2 Potatoes Chemicals and Seed-borne Fungi

### Rothamsted Research

Rothamsted Research (1971) *70/R/P/2 Potatoes Chemicals and Seed-borne Fungi* ; Numerical Results Of The Field Experiments 1970, pp 293 - 294 - DOI:

<https://doi.org/10.23637/ERADOC-1-59>

POTATOES

(70/R/P/2)

Chemicals and seed-borne fungi, Great Knott II 1970.

Design: 4 randomised blocks of 2 plots, split into 8. (Plus one extra block for sampling).

Area of each sub-plot: 0.0036. Area harvested: 0.0036.

Treatments: All combinations of:-

Whole plots: 1. Varieties: King Edward (KE), Pentland Crown (PC).

Sub-plots: 2. Seed treatment with chemicals: Benomyl powder (B), thiabendazole powder (T).

3. Rates of application: Diluted with kaolin dust to give 1% (1), 5% (2), 10% (3) a.i.

Together with additional treatments to sub-plots: None (0), kaolin dust (K).

NOTE: (1) All dust treatments were at 10 lb per ton of seed potatoes.

Basal applications: 10 cwt (13:13:20). Weedkiller: Linuron at 1 lb plus paraquat at 0.375 lb ion in 39 gals. Fungicide: Mancozeb at 1.2 lb in 40 gals on 2 occasions. Insecticide: Demeton-s-methyl at 3.5 oz applied with the first mancozeb spray.

Cultivations, etc.: Ploughed: 28 Oct, 1969. Basal NPK applied: 23 Apr, 1970. Rotary cultivated: 7 May. Potatoes machine planted: 8 May. Weedkiller applied: 23 May. Fungicide and insecticide applied: 29 July. Fungicide applied: 14 Aug. Sprayed with undiluted BOV at 15 gals: 3 Sept. Lifted: 18 Sept. Previous crops: Barley 1968, fallow 1969.

NOTE: (2) Emergence counts were made, and samples were taken during the season for disease assessments. Diseases in the eyes of tubers were assessed after riddling.

Standard errors per plot. Total tubers, tons:

Whole plot: 1.004 or 7.6% (3 d.f.)

Sub plot: 0.578 or 4.4% (42 d.f.)

70/R/P/2

SUMMARY OF RESULTS

	O	K	B1	B2	B3	T1	T2	T3	Mean
TOTAL TUBERS: TONS									
	(1) and (2)								(±0.502)
KE	13.33	13.37	13.87	13.44	13.71	14.10	13.59	13.59	13.62
PC	12.69	12.91	13.24	12.86	12.95	12.42	12.43	12.82	12.79
Mean (±0.205)	13.01	13.14	13.55	13.15	13.33	13.26	13.01	13.20	13.21

(1) (±0.570) For use in vertical and diagonal comparisons only  
 (2) (±0.289) For use in horizontal and interaction comparisons only

¾ WARE: 1.75 INCH

KE	74.5	77.6	76.9	75.6	76.2	78.5	78.3	79.2	77.1
PC	95.6	95.9	95.8	96.4	96.0	95.4	95.9	95.2	95.8
Mean	85.0	86.7	86.4	86.0	86.1	87.0	87.1	87.2	86.4