

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/P/6 Potatoes Systemic Nematicide

Rothamsted Research

Rothamsted Research (1971) *70/W/P/6 Potatoes Systemic Nematicide* ; Numerical Results Of The Field Experiments 1970, pp 304 - 307 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

POTATOES

(70/W/P/6)

Systemic nematicide - Woburn Butt Close 1970.

Design: 3 blocks of 8 plots.

Area of each plot: 0.0026. Area harvested: 0.0013.

Treatments: Nematicide:

Broadcast: None (S0), 4.3 (S1), 8.6 (S2), 17.2 lb (S4) of 10% granules of DP 1410 (S-methyl 1-(dimethylcarbamoyl)-N-((methylcarbamoyl)oxy)thioformimidate).

Broadcast and Foliar spray: 4.3 + 2 lb (S1F1), 4.3 + 4 lb (S1F2), 8.6 + 2 lb (S2F1), 8.6 + 4 lb (S2F2) of DP 1410.

Basal applications: 10 cwt (13:13:20) broadcast, plus 7.5 cwt (13:13:20) placed. Weedkiller: Linuron at 1 lb in 33 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on two occasions. Insecticide: Demeton-s-methyl at 3.5 fl oz on one occasion with fungicide.

Cultivations, etc.: Deep-tine cultivated: 3 Nov, 1969. NPK broadcast, deep-tine cultivated, 2 Apr, 1970. Nematicide broadcast: 4 May. NPK placed, potatoes planted, inter-row rotary cultivated and earthed up: 5 May. Weedkiller applied: 15 May. Nematicide spray applied: 3 June. Fungicide applied: 31 July, 18 Aug. Insecticide applied: 31 July. Lifted: 29 Sept. Variety: Pentland Crown. Previous crops: Potatoes 1968, 1969.

NOTE: Soil samples were taken in May and after harvest for egg and larvae counts of *Heterodera rostochiensis*.

Standard error per plot.

Total tubers, tons: 1.413 or 8.4% (14 d.f.)

70/W/P/6

SUMMARY OF RESULTS

S0	S1	S2	S4	S1F1	S1F2	S2F1	S2F2	Mean
TOTAL TUBERS: TONS								
(±0.816)								
9.43	18.23	19.45	19.04	16.38	16.84	18.40	17.59	16.92
% WARE: 1.5 INCH								
89.9	92.9	94.0	92.7	93.3	93.4	94.3	91.5	92.8

POTATOES

(70/R/P/11)

Varieties and 'Ethrel', Long Hoos IV 1970.

Design: 4 blocks of 8 plots, randomisation restricted.

Area of each plot: 0.0051. Area harvested: 0.0006.

Treatments: All combinations of:-

1. Varieties: Arran Pilot (AP), King Edward (KE).
2. 'Ethrel':* Seed soaked for 1 hour before planting: Not soaked (E0), 'Ethrel' at 60 p.p.m. (E1), 120 p.p.m. (E2), 240 p.p.m. (E4).

* 2-chloroethylphosphonic acid.

Basal applications: 1120 lb (13:13:20). Weedkiller: Linuron at 1 lb in 39 gals. Fungicide: Mancozeb at 1.2 lb in 40 gals on 2 occasions. Insecticide: Demeton-s-methyl at 3.5 oz with first mancozeb spray.

Cultivations, etc.: Ploughed: 28 Oct, 1969. Basal NPK applied: 23 Apr, 1970. Plots rotary cultivated, potatoes machine planted: 30 Apr. Weedkiller applied: 23 May. Grubbed and rotary ridged: 17 June. Fungicide and insecticide applied: 29 July. Fungicide applied: 14 Aug. Lifted: 9 Sept. Previous crops: Barley 1968, spring beans 1969.

NOTE: Samples were taken every month for estimation of leaf areas, tuber numbers and tuber size distribution.

Standard error per plot.

Total tubers, tons: 1.224 or 11.9% (21 d.f.)

70/R/P/11

SUMMARY OF RESULTS

	EO	E1	E2	E4	Mean
TOTAL TUBERS: TONS					
(±0.612)					
AP	7.95	9.60	7.91	7.65	8.28
KE	11.92	12.05	12.42	12.65	12.26
Mean (±0.433)	9.93	10.83	10.17	10.15	10.27

% WARE: 1.5 INCH

AP	90.6	94.2	94.0	92.9	92.9
KE	94.7	94.2	92.7	94.9	94.1
Mean	92.7	94.2	93.3	93.9	93.5