Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1970



Full Table of Content

70/R/G/3 Grass Weedkiller and Aqueous N

Rothamsted Research

Rothamsted Research (1971) 70/R/G/3 Grass Weedkiller and Aqueous N; Numerical Results Of The Field Experiments 1970, pp 315 - 319 - DOI: https://doi.org/10.23637/ERADOC-1-59

GRASS

(70/R/G/3)

Weedkiller and aqueous N, Road Piece 1970, permanent grass.

Design: 4 randomised blocks of 28 plots.

Area of each plot: 0.0007. Area harvested: 0.0004.

Treatments: All combinations of:-

- 1. Weedkiller (dichlorprop/MCPA): None (HO), 20 (H1), 40 (H2), 60 (H3) oz total a.e. on 2 occasions.
- 2. Forms of nitrogen (on 2 occasions): Solid, as 'Nitro-Chalk' (21% N) applied immediately after the weedkiller (S), liquid as urea/ammonium nitrate (26% N) mixed with weedkiller (L).
- 3. Levels of nitrogen: 0.3 (N1), 0.6 (N2), 0.9 (N3) cwt N on each occasion.

Together with 4 additional treatments:

SN2 E HD, SN2 E H1, SN2 E H2, SN2 E H3

where 'Nitro-Chalk' was applied early (E) and the HD plots were treated with weedkiller on 2 occasions (Dalapon at 4 lb a.i. in 28 gals).

NOTE: The dichlorprop/MCPA was applied in 28 gals where solid fertiliser was used. The liquid fertiliser (with or without weedkiller) was applied as a spray in 10, 20 and 30 gals for rates 1, 2 and 3 respectively.

Basal applications: 560 lb (0:14:28).

- Cultivations, etc.: Basal PK applied: 10 Apr, 1970. N applied to E plots: 27 Apr. Remaining N treatments and weedkillers (including dalapon) applied: 19 May. Cut: 16 June. N applied to E plots: 22 June. Remaining N treatments and weedkillers (including dalapon) applied: 30 June. Cut second time: 15 Oct.
- NOTES: (1) Soil samples were taken for pH in May. Scores were made of weedkiller scorch and weed control. Scores were also made for grass cover and for fine-leaved grasses. On blocks II and IV the proportion of weeds to grasses at the second cut was determined.

70/R/G/3

- (2) On the second cut on one plot L N3 HO the material harvested was mainly docks. An estimated value was used in the analysis for second cut and total of 2 cuts.
- (3) The percentage of N in herbage was determined.

Standard errors per plot. Dry matter: cwt.

1st cut:
3.92 or 16.4% (69 d.f.)
2nd cut:
3.65 or 14.0% (68 d.f.)
Total of 2 cuts: 5.09 or 10.2% (68 d.f.)

70/R/G/3 SUMMARY OF RESULTS

DRY MATTER: CWT

	NI	N2	N3	HO	H1	H2	Н3	Mean	
72			1ST	CUT	178 6	30-			
	(±0.98)				(±0.57)				
S L			23.5	26.5 25.1	25.5 24.9	21.7	22.0		
					(±1.38)				
		N3 N2 N1		27.1	27.6 24.0 24.1	23.2	22.7	24.3	
	23 13	Mean (±0.80)		25.8	25.2	22.3	22.0	23.8	
		SN2 E SN2 E SN2 E SN2 E	H1 H2	18.5 22.9 26.9 25.5					

General mean: 23.8 Mean D.M. %: 30.9

70/R/G/3

DRY MATTER: CWT

	Nl	N2	N3	НО	Hl	H2	Н3	Mean	
	2ND CUT								
		(±0.91)			(±1.05)				
3	26.4	27.2	30.2	30.4	27.1 23.3	27.7	26.6	27.9 24.1	
					(±1.29)				
		N1 N2 N3		28.4	24.2	22.4 26.5 28.8	24.8	26.0	
	-63	Mean (±0.7		28.1	25.2	25.9	24.9	26.0	
			H1 H2	8.6 28.5 24.1 28.1					

General mean: 25.5 Mean D.M.%: 26.4

70/R/G/3
DRY MATTER: CWT

	Nl	N2	M3	OH	H1	H2	Н3	Mean		
	TOTAL OF 2 CUTS									
	(±1.27)				(±0.73)					
S L	50.4 44.8	51.5 49.0	53.7 49.5	56.8 50.6	52.7 48.2	49.4 47.0	48.6 45.1	5 1. 9 47.8		
					(±0.90)					
	N1 N2 N3 Mean (±1.04)			55.5	51.0 48.2 52.2	49.8	47.5	50.2		
				53.7	50.5	48.2	46.9	49.8		
		SN2 E SN2 E SN2 E	H1 H2	27.1 51.3 51.1 53.7						

General mean: 49.2 Mean D.M. %: