Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



## Numerical Results of the Field Experiments 1970



Full Table of Content

## 70/G/SC/1 Sweet Corn Nitrogen and Seed Rates

## **Rothamsted Research**

Rothamsted Research (1971) 70/G/SC/1 Sweet Corn Nitrogen and Seed Rates; Numerical Results Of The Field Experiments 1970, pp 320 - 321 - DOI: https://doi.org/10.23637/ERADOC-1-59

SWEET CORN

(70/G/SC/1)

Nitrogen and seed rates Rothamsted Garden Plots.

Design: 4 randomised blocks of 12 plots.

Area of each plot: 0.0031. Area harvested: 0.0012.

Treatments: All combinations of:-

- Seed spacing: 3 in. (R1), 6 in. (R2), 12 in. (R3) in rows 2 feet apart.
- Nitrogen: None (NO), 0.5 (N1), 1.0 (N2), 1.5 (N3) cwt as 'Nitro-Chalk'.
- Basal applications: 1230 lb (0:14:28) broadcast. Insecticide:
  Phorate at 1.5 lb drilled at 4 in. depth in the rows. Weedkiller:
  Atrazine at 1.5 lb in 35 gals.
- Cultivations, etc.: Basal PK applied: 21 Nov, 1969. Ploughed: 23 Dec. Insecticide applied, seed drilled: 15 May, 1970. Weedkiller applied: 20 May. N applied: 3 June. Harvested by hand: 9 Sept. Variety: Early King. Previous crops: Fallow 1968, potatoes 1969.

NOTE: Cobs were counted and weighed in 2 grades:

1. Large saleable, 2. Small saleable.

Standard errors per plot:

Total saleable cobs, cwt:

Number of saleable cobs, thousands:

2.74 or 13.1 (33 d.f.)

70/G/SC/1

## SUMMARY OF RESULTS

	NO	Nl	N2	N3	Mean
	TO	TAL SALEABL	E COBS: CWT	Terrory.	
1 1000	(±5.84)				(±2,92)
R1 R2 R3	88.6 86.6 78.0	112.8 92.0 78.7	138.7 108.1 97.4	118.2 107.3 80.6	114.6 98.5 83.7
Mean (±3.37)	84.4	94.5	114.7	102.0	98.9
	NUMBER	OF SALEABLE	COBS: THOUSA	ANDS	
	(±1.37)				
R1 R2 R3	20.5 18.6 16.1	24.9 18.4 16.8	31.8 21.4 19.5	25.8 21.2 16.1	25.8 19.9 17.1
Mean (±0.79)	18.4	20.0	24.2	21.0	20.9