

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)

International Agricultural Research
Institute
London Agricultural Station
Harpenden, Herts
AL5 2JQ
ENGLAND
01582 555000
© 2015 Rothamsted Research. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without the prior written permission of Rothamsted Research. For more information, please contact the Rothamsted Research Library, Rothamsted, Harpenden, Herts, AL5 2JQ, UK. Email: library@rothamsted.ac.uk

70/R/M/4 and 70/W/M/4 Mixed Crops Ammonium Phosphates for Grass and Barley

Rothamsted Research

Rothamsted Research (1971) *70/R/M/4 and 70/W/M/4 Mixed Crops Ammonium Phosphates for Grass and Barley* ; Numerical Results Of The Field Experiments 1970, pp 322 - 329 - DOI:
<https://doi.org/10.23637/ERADOC-1-59>

MIXED CROPS

(70/R/M/4 and 70/W/M/4)

Ammonium phosphates for grass and barley, Rothamsted (R) West Barnfield II and Woburn (W) Stackyard Series B sideland, 1970.

Design (each field):-

Ryegrass: 3 randomised blocks of 16 plots.
Barley: 3 randomised blocks of 12 plots.

Area of each plot:

Ryegrass: 0.0010. Area harvested: 0.0006.
Barley: 0.0003. Area harvested: 0.00003.

Treatments:

No fertiliser	(O)
Mono-ammonium phosphate + ammonium sulphate*	(A)
Mono-sodium phosphoramidate + ammonium sulphate*	(M)
Sodium phosphordiamidate + ammonium sulphate*	(D)
Phosphoryl triamide	(T)

* To give the same N:P ratio as treatment T.

Ammonium nitrate to grass: 200 lb N in seedbed	(N)
Ammonium nitrate to barley: 100 lb N West Barnfield II (R), 150 lb N Stackyard B (W)	(N)

The treatments described above were applied either singly or in combinations, as shown below:-

To ryegrass: O (4 plots per block) and all combinations of:-

1. NP fertilisers: A, M, D, T.
2. Rates of fertiliser: To give 10 lb P, 13.5 lb N from NP fertiliser, plus ammonium nitrate at rate given above (1+N), to give 20 lb P, 27 lb N from NP fertiliser, plus ammonium nitrate (2+N), to give 20 lb P, 27 lb N from NP fertiliser (2).

To barley: O, N, monosodium phosphate at 10 lb P (P1) and 20 lb (P2) and all combinations of:-

1. NP fertilisers: A, M, D, T.
2. Rates of fertiliser: To give 10 lb P, 13.5 lb N from NP fertiliser, plus ammonium nitrate at rate given above (1+N), to give 20 lb P, 27 lb N from NP fertiliser, plus ammonium nitrate (2+N).

70/R&W/M/4

Basal applications:

Ryegrass: 100 lb K as potassium sulphate. Weedkiller: Mecoprop ('Clovotox' at 4 pints in 45 gals).
Barley: 50 lb K as potassium sulphate.

Cultivations, etc.:-

West Barnfield II (R): Ploughed: 29 Sept, 1969.

Ryegrass: Treatment fertilisers and basal K applied: 2 Apr, 1970.
Seed drilled at 30 lb: 17 Apr. Weedkiller applied: 29 May.
Cut twice: 5 Aug and 16 Sept. Variety: S22 Italian.

Barley: Treatment fertilisers and basal K applied: 2 Apr, 1970.
Seed drilled at 150 lb: 16 Apr. Cut green: 10 and 29 June (separate areas). Variety: Julia. Previous crops: Barley 1968 and 1969.

Stackyard B (W): Ploughed: 19 Nov, 1969.

Ryegrass: Treatment fertilisers and basal K applied: 3 Apr, 1970.
Seed drilled at 30 lb: 7 Apr. Weedkiller applied: 29 May.
Cut twice: 22 July and 17 Sept. Variety: S22 Italian.

Barley: Treatment fertilisers and basal K applied: 3 Apr, 1970.
Seed drilled at 150 lb: 7 Apr. Cut green: 3 and 24 June (separate areas). Variety: Julia. Previous crops: Fallow 1968 and 1969.

NOTE: Crop samples were taken for determination of dry matter and N %.

Standard errors per plot.

Ryegrass, dry matter, cwt:

West Barnfield II (R): 1st cut: 1.36 or 13.5% (22 d.f.)
2nd cut: 1.44 or 8.7% (22 d.f.)
Total of 2 cuts: 2.08 or 7.9% (22 d.f.)

Stackyard B (W): 1st cut: 2.08 or 19.2% (22 d.f.)
2nd cut: 2.20 or 13.0% (22 d.f.)
Total of 2 cuts: 3.62 or 13.0% (22 d.f.)

Barley, green crop, dry matter, cwt:

West Barnfield II (R): 1st cut: 2.53 or 19.2% (18 d.f.)
2nd cut: 5.88 or 15.7% (18 d.f.)

Stackyard B (W): 1st cut: 5.17 or 21.0% (18 d.f.)
2nd cut: 6.04 or 10.5% (18 d.f.)

70/R&W/M/4

SUMMARY OF RESULTS

RYEGRASS

DRY MATTER: CWT

WEST BARNFIELD II (R)

	A	M	D	T	Mean
1ST CUT					
(±0.78)					
1+N	12.1	9.7	10.8	11.3	11.0
2+N	9.7	11.1	13.3	10.3	11.1
2	7.8	8.3	9.0	7.2	8.1
Mean (±0.45)	9.8	9.7	11.0	9.6	10.0

O: 5.5

General mean: 8.9

Mean D.M. %: 35.1

2ND CUT					
(±0.83)					
1+N	17.6	19.4	19.3	18.0	18.6
2+N	16.9	18.1	18.5	17.7	17.8
2	13.3	12.9	13.4	12.3	13.0
Mean (±0.48)	15.9	16.8	17.1	16.0	16.5

O: 10.4

General mean: 14.9

Mean D.M. %: 23.0

70/R&N/M/4

RYEGRASS

DRY MATTER: CWT

WEST BARNFIELD II (R)

TOTAL OF 2 CUTS

	A	M	D	T	Mean
		(±1.20)			(±0.60)
1+N	29.7	29.1	30.1	29.3	29.6
2+N	26.6	29.2	31.8	28.0	28.9
2	21.1	21.2	22.4	19.5	21.0
Mean (±0.69)	25.8	26.5	28.1	25.6	26.5

O: 15.9

General mean: 23.9

Mean D.M. %: 29.0

70/R&W/M/4

RYEGRASS

DRY MATTER: CWT

STACKYARD B (W)

	A	M	D	T	Mean
1ST CUT					
(±1.20)					
1+N	15.2	14.4	12.5	14.7	14.2
2+N	12.5	13.6	15.1	14.6	13.9
2	4.5	4.1	5.4	3.6	4.4
Mean (±0.69)	10.7	10.7	11.0	11.0	10.8

O: 1.1

General mean: 8.4

Mean D.M. %: 31.1

	A	M	D	T	Mean
2ND CUT					
(±1.27)					
1+N	22.0	20.2	19.2	23.6	21.2
2+N	21.5	21.3	21.7	22.3	21.7
2	7.5	8.4	9.1	6.9	8.0
Mean (±0.73)	17.0	16.6	16.7	17.6	17.0

O: 5.1

General mean: 14.0

Mean D.M. %: 32.8

70/R&W/W/4

RYEGRASS

DRY MATTER: CWT

STACKYARD B (W)

TOTAL OF 2 CUTS

	A	M	D	T	Mean
			(±2.09)		(±1.04)
1+N	37.2	34.6	31.7	38.3	35.4
2+N	34.0	34.9	36.8	36.9	35.6
2	12.0	12.6	14.5	10.5	12.4
Mean (±1.21)	27.7	27.4	27.7	28.6	27.8

O: 6.3

General mean: 22.4

Mean D.M. %: 32.0

70/R&N/M/4

BARLEY

GREEN CROP, DRY MATTER: CWT

WEST BARNFIELD II (R)

	P*	A	M	D	T	Mean
1ST CUT						
	(±1.46)					(±0.65)
1+N	9.9	10.9	12.2	13.7	14.0	12.1
2+N	10.9	16.5	13.3	16.1	14.7	14.3
Mean (±1.03)	10.4	13.7	12.7	14.9	14.4	13.2
O:	6.5					
N:	10.8					
General mean:	12.5					
2ND CUT						
	(±3.39)					(±1.52)
1+N	28.2	37.7	38.5	42.8	37.2	36.9
2+N	30.7	39.4	45.1	37.6	37.7	38.1
Mean (±2.40)	29.5	38.6	41.8	40.2	37.5	37.5
O:	22.8					
N:	31.4					
General mean:	35.8					

* Did not receive N

70/R&M/4

BARLEY

GREEN CROP, DRY MATTER: CWT

STACKYARD B (W)

	P*	A	M	D	T	Mean
1ST CUT						
			(±2.99)			(±1.34)
1+N	3.9	25.2	30.1	27.8	29.1	23.2
2+N	4.5	36.3	28.9	30.8	29.3	26.0
Mean (±2.11)	4.2	30.8	29.5	29.3	29.2	24.6
C:	4.9					
N:	27.0					
General mean:	23.2					
2ND CUT						
			(±3.49)			(±1.56)
1+N	13.3	67.6	72.0	69.6	62.7	57.0
2+N	12.0	68.6	72.0	64.9	71.1	57.7
Mean (±2.47)	12.7	68.1	72.0	67.2	66.9	57.4
C:	13.6					
N:	66.8					
General mean:	53.6					
* Did not receive N						