

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/RN/7 Residual Phosphate - Potatoes, Barley, Turnips

Rothamsted Research

Rothamsted Research (1971) *70/R/RN/7 Residual Phosphate - Potatoes, Barley, Turnips* ; Numerical Results Of The Field Experiments 1970, pp 90 - 93 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

RESIDUAL PHOSPHATE

(70/R/RN/7)

Potatoes, barley and swedes, Great Field IV and Sawyers I the 11th year. For treatments and previous years' results see 'Details' 1967 and 'Results' 67/B/6, 68/B/5 and 69/R/RN/7.

Area of each plot:-

Great Field IV: 0.0193. Area harvested: Potatoes and barley - 0.0129.
Turnips - 0.0096.

Sawyers I: 0.0212. Area harvested: Potatoes and barley - 0.0141.

Revised manuring on both fields:

	cwt N	cwt K ₂ O
Potatoes	2.0	2.0 as 'Nitro-Chalk' and sulphate of potash
Barley	0.8	0.5 as (25:0:16) combine drilled
Swedes	0.5	1.0 as 'Nitro-Chalk' and sulphate of potash

The barley variety is now Julia.

NOTE: Because of dry weather, swedes failed on both fields and turnips were sown (without further manuring). On Sawyers I the turnips were seriously affected by *Phoma* and yields were not taken.

Standard applications:

Potatoes: Weedkiller: Linuron at 1 lb plus paraquat at 0.375 lb ion in 39 gals. Fungicide: Mancozeb at 1.2 lb in 40 gals on 2 occasions. Insecticide: Demeton-s-methyl at 3.5 oz with first mancozeb spray.

Barley: Ground chalk at 25 cwt. Weedkiller: 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc. (both fields):- Ploughed: 24 Nov, 1969.

Potatoes: Fertilisers applied: 5 May, 1970. Plots rotary cultivated, potatoes machine planted: 6 May. Weedkiller applied: 23 May. Grubbed and rotary ridged: 18 June. Fungicide and insecticide applied: 29 July. Fungicide applied: 14 Aug. Sprayed with undiluted BOV at 15 gals: 15 Sept. Lifted: 25 Sept.

Barley: Ground chalk applied: 6 Oct, 1969. Treatment P fertilisers applied: 26 Mar, 1970. Seed combine drilled at 140 lb: 28 Mar. Weedkiller applied 18 May. Combine harvested: 14 Aug. Variety: Julia.

70/R/RN/7

Swedes: Fertilisers applied: 14 May, 1970. Seed drilled at 1.5 lb: 15 May. Power harrowed for turnips: 23 June. Rotary cultivated: 24 June. Turnips drilled at 1.5 lb: 25 June. Singled: 23 July - 7 Aug. Lifted: 12 Oct. Variety of turnips: Imperial Green Globe.

Standard errors per plot.

Sawyers I:

Potatoes, total tubers, tons: 1.016 or 9.5% (11 d.f.)

Barley, grain, cwt: 1.57 or 5.1% (11 d.f.)

70/R/RN/7

SUMMARY OF RESULTS

POTATOES

Treatment	TOTAL TUBERS: TONS		% WARE: 1.5 INCH	
	Great Field IV Mean	Sawyers I Mean	Great Field IV Mean	Sawyers I Mean
		(±0.719)		
O	11.68	9.25	98.0	98.2
A1	12.87	8.54	98.5	96.4
A2	15.32	10.12	98.2	98.1
A3	16.80	11.86	98.5	96.9
A4	17.32	13.44	98.1	97.9
T1	12.18	8.43	98.4	97.3
T2	14.17	9.89	97.5	97.1
R2	15.21	12.50	98.3	98.3
R3	14.37	11.51	97.4	97.8
R4	15.53	13.70	97.3	97.5
G1	12.62	7.60	97.6	98.0
S1	13.55	11.00	98.3	98.1
Mean	14.30	10.65	98.0	97.6

BARLEY

	GRAIN: CWT		STRAW: CWT	
		(±1.11)		
O	23.9	24.1	16.9	8.6
A1	36.6	30.5	17.7	10.5
A2	37.8	31.9	18.9	11.5
A3	33.9	33.2	19.7	10.8
A4	33.7	34.3	24.7	14.3
T1	42.0	28.0	17.3	9.4
T2	36.4	30.9	16.3	11.2
R2	32.1	31.7	14.4	8.9
R3	32.8	33.0	19.8	11.5
R4	33.1	33.7	19.5	13.2
G1	36.5	28.6	18.1	9.5
S1	32.3	27.6	15.1	9.8
Mean	34.3	30.6	18.2	10.8
Mean D.M.%:	86.5	86.7	88.2	88.1

70/R/RIV/7

TURNIPS, ROOTS: TONS

Treat- ment	Great Field IV Mean
O	0.37
A1	4.40
A2	6.76
A3	8.47
A4	6.95
T1	3.15
T2	3.38
R2	3.98
R3	4.82
R4	5.37
G1	2.69
S1	2.32
Mean	4.39