

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



### 70/R/RN/9 Cereal Disease Reference Plots - W. Wheat, S. Wheat, Oats, S. Beans

#### Rothamsted Research

Rothamsted Research (1971) *70/R/RN/9 Cereal Disease Reference Plots - W. Wheat, S. Wheat, Oats, S. Beans* ; Numerical Results Of The Field Experiments 1970, pp 101 - 102 - **DOI:**  
<https://doi.org/10.23637/ERADOC-1-59>

CEREAL DISEASE REFERENCE PLOTS

(70/R/RN/9)

Pennells Piece 1970, the eighth year, winter and spring wheat.  
For treatments etc., see 'Results' 63/C/10 (W = Wheat,  
O = Oats, BE = Spring beans).

Area of each plot: 0.0180. Area harvested: 0.0118.

Cultivations, etc.: Sprayed with paraquat at 0.5 lb ion in 20 gals:  
18 Sept, 1969. Ploughed: 19 Sept.

Winter wheat: Rotary cultivated: 15 Oct, 1969. Seed combine  
drilled at 180 lb, plots sprayed with terbutryne and related  
triazines ('Prebane' at 4 lb in 20 gals): 17 Oct. N applied:  
25 Mar, 1970. Sprayed with 2,4-D at 0.5 lb and dichlorprop  
at 2 lb in 20 gals: 13 May. Combine harvested: 26 Aug.

Spring wheat: Seed combine drilled at 170 lb: 22 Apr, 1970.  
N applied: 24 Apr. Sprayed with ioxynil at 7.5 oz and  
mecoprop at 22.5 oz in 20 gals: 27 May. Combine harvested:  
28 Aug.

Oats: Seed combine drilled at 170 lb: 17 Apr, 1970. N applied:  
24 April. Combine harvested: 15 Aug.

Spring beans: Seed placement drilled at 200 lb: 21 Mar, 1970.  
Phorate at 17 oz in granules applied: 16 June. Combine  
harvested: 4 Sept.

- NOTES: (1) Yields were taken for winter and spring wheat only-  
(Crop sequences 1, 4, 5 and 6).  
(2) Estimates were made in spring and summer of the  
incidence of take-all (*Ophiobolus graminis*) and  
eyespot (*Cercospora herpotrichoides*).  
(3) For previous years' results see 'Results' 63/C/10,  
64/C/9, 65/C/9, 66/C/7, 67/C/5, 68/C/5 and 69/R/RN/9.

70/R/RN/9

SUMMARY OF RESULTS

GRAIN: CWT

Crop in	C1	C4	C5	C6	
1963	W	BE	O	W	
1964	W	O	W	W	
1965	W	W	W	W	
1966	BE	W	W	W	
1967	O	W	BE	W	
1968	W	BE	O	W	
1969	W	O	W	W	Mean

WINTER WHEAT

42.0	43.7	44.4	39.8	42.5
------	------	------	------	------

SPRING WHEAT

16.7	23.8	22.1	19.2	20.4
------	------	------	------	------

Mean D.M. %: Winter wheat: 83.9  
 Spring wheat: 84.2