


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/W/RN/12 Organic Manuring - Leys, Rye

Rothamsted Research

Rothamsted Research (1972) 71/W/RN/12 *Organic Manuring - Leys, Rye ; Yields Of The Field Experiments 1971*, pp 107 - 110 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/W/RN/12

ORGANIC MANURING

Object: To study, from crop yields and soil analyses, the cumulative effects of a range of types of organic matter - Woburn Stackyard B.

The seventh year, leys and rye.

For previous years see 66/C/31(t), 67/C/24(t), 68/C/18(t), 69/W/RN/12(t) and 70/W/RN/12(t).

Whole plot dimensions: 8.53 x 30.5. Area harvested: Leys - 0.00524, rye - 0.00421.

Fertilisers applied autumn 1970 (kg)

Treatment	P205	K20	MgO
DG	-	-	-
ST	63	-	25
PT	63	126	-
GM	63	138	38
FD	113	427	38
FS	75	126	38
LC	63	126	38
LN	63	126	38

Fertilisers applied spring 1971 (kg)

Treatment	P205	K20
ST	-	50
LC	63	126
LN	88	251

No P, K, or Mg was required in spring for DG, PT, GM, FD or FS.

NOTE: Residues of N to sugar beet 1969
(N1), (N3), (N5), (N7). 25, 75, 125, 175 kg N as 'Nitro-Chalk' 21.

Basal applications: Rye: 31 kg N as 'Nitro-Chalk' 21.
LN plots: 126 kg N as 'Nitro-Chalk' 21 in spring and after each cut except the last.

Cultivations, etc.:

LC and LN plots: P, K, Mg applied: 6 Oct, 1970. P, K applied: 9 Mar, 1971. N applied to LN plots: 16 Mar, 29 June. Cut: 23 June, 10 Sept.

Rye: P, K, Mg applied: 17 - 18 Sept, 1970. Peat, straw applied: 15 Oct. FYM applied, ploughed: 16 - 17 Oct. Seed drilled

71/W/RN/12

at 191 kg: 17 Oct. GM plots undersown with Late Flowering Red Clover at 72 kg: 19 Oct. K applied to ST plots only: 9 Mar, 1971. N applied: 14 Apr. Red Clover resown at 72 kg: 22 Apr. Combine harvested: 31 Aug. Variety: King II.

NOTE: Soil samples were taken in the stubble for P, K and Mg analysis.

Standard errors per plot. Rye, grain, tonnes/hectare:
Whole plot: 0.266 or 7.3% (15 d.f.)
Sub plot: 0.199 or 5.5% (54 d.f.)

71/W/RN/12

SUMMARY OF RESULTS

LEYS

DRY MATTER: TONNES/HECTARE

LC		LN
<hr/>		
	1ST CUT	
2.15		6.51
2ND CUT		
2.13		4.40
TOTAL OF 2 CUTS		
4.28		10.91

Mean D.M. %: 1st cut: 24.6
2nd cut: 24.1
Total of 2 cuts: 24.4

71/W/RN/12

RYE

1969

	N1	N3	N5	N7	Mean
GRAIN: TONNES/HECTARE					
	(1) and (2)				
DG	4.76	4.63	4.46	4.63	4.62
ST	3.61	3.63	3.48	3.76	3.62
PT	3.48	3.43	3.48	3.58	3.49
GM	3.33	3.18	3.46	3.44	3.35
FD	3.29	3.38	3.48	3.32	3.37
FS	3.33	3.46	3.36	3.03	3.29
Mean (±0.041)	3.63	3.62	3.62	3.63	3.62

(1) (± 0.158) For use in vertical and diagonal comparisons only

(2) (± 0.099) For use in horizontal and interaction comparisons only

	STRAW: TONNES/HECTARE				
DG	6.27	6.34	5.68	6.11	6.10
ST	4.53	4.46	4.18	4.56	4.43
PT	4.40	4.26	4.40	4.33	4.35
GM	4.25	4.70	5.20	4.53	4.67
FD	4.22	4.41	4.32	4.48	4.36
FS	4.05	4.09	4.05	4.05	4.06
Mean	4.62	4.71	4.64	4.67	4.66

Mean D.M. %: Grain: 75.1
Straw: 88.0