


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1971

[Full Table of Content](#)



71/W/RN/13 Intensive Cereals - Ley, Potatoes, Wheat, Barley

Rothamsted Research

Rothamsted Research (1972) 71/W/RN/13 *Intensive Cereals - Ley, Potatoes, Wheat, Barley* ; Yields Of The Field Experiments 1971, pp 111 - 115 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne diseases and organic matter in the soil-Woburn Stackyard I.

The sixth year, ley, potatoes, winter wheat, barley.

For previous years see 66/B/9(t), 67/B/9, 68/B/7(t) and 69-70/W/RN/13(t).

Design: For each cereal: 2 randomised blocks of 6 plots, split into 4.

Whole plot dimensions: 8.53 x 20.4. Sub plot area harvested:
Potatoes - 0.00343, wheat - 0.00271, barley - 0.00277.

NOTE: The magnesium test on all crops is now ended.

Basal applications: All crops: 5020 kg magnesian limestone three-quarters ploughed in, a quarter applied to plough furrow. 126 kg P2O5, 252 kg K2O as (0:14:28), half ploughed in, half applied to plough furrow.

Weedkiller: Paraquat at 1.12 kg ion in 281 l.

Ley: 63 kg N in seedbed, 63 kg N 8 weeks after sowing, 75 kg N after each cut except the last as 'Nitro-Chalk' 21.

Potatoes: 150 kg N as 'Nitro-Chalk' 21. Weedkiller: Linuron at 1.12 kg in 371 l. Fungicide: Mancozeb at 1.34 kg in 416 l on the first occasion, and 371 l on the second. Insecticide:

Demeton-s-methyl at 246 ml applied with first mancozeb spray.

Wheat and barley: Weedkiller: Ioxynil at 0.53 Kg and mecoprop at 1.58 kg in 281 l.

Cultivations, etc.: All plots: Paraquat applied: 16 Sept, 1970. Half PK applied: 24 Sept. Three-quarters magnesian limestone applied: 6 Oct. Ploughed: 10 Oct. Remaining PK and magnesian limestone applied: 14 Oct.

Leys: Deep-tine cultivated barley blocks: 21 Oct, 1970. N applied: 1 Apr. Seeds sown at 33 kg: 22 Apr. First cut: 16 July. N applied: 19 July. Second cut: 8 Sept. Variety: Italian Ryegrass S22.

Potatoes: Deep-tine cultivated barley blocks: 21 Oct, 1970. N applied: 29 Mar, 1971. Rotary cultivated, potatoes planted: 30 Mar. Weedkiller applied: 5 May. Rotary ridged: 1 June.

Fungicide plus insecticide applied: 28 June. Fungicide applied: 12 Aug. Haulm mechanically destroyed: 13 Sept. Lifted: 14 Sept. Variety: Majestic.

Wheat: Seed drilled at 191 kg: 14 Oct, 1970. N applied: 14 Apr, 1971. Weedkiller applied: 28 Apr. Combine harvested: 28 Aug. Variety: Cappelle.

71/W/RN/13

Barley: Deep-tine cultivated: 21 Oct, 1970. N applied, seed drilled at 157 kg: 17 Mar, 1971. Power harrowed, re-drilled at 157 kg: 9 Apr. Weedkiller applied: 18 May. Combine harvested: 25 Aug. Variety: Julia.

NOTE: Estimates of eyespot (*Cercosporaella herpotrichoides*) and take-all (*Ophiobolus graminis*) were made on both cereal crops.

Standard errors per sub plot.

Wheat, grain, tonnes/hectare: Sub plot: 0.391 or 10.6% (12 d.f.)
Barley, grain, tonnes/hectare: Sub plot: 0.194 or 4.5% (11 d.f.)

71/W/RN/13

SUMMARY OF RESULTS

POTATOES

N: KG/HA 1969

63	126	189	252	Mean
----	-----	-----	-----	------

PERMANENT WHEAT BLOCKS

TOTAL TUBERS: TONNES/HECTARE

33.0	32.2	36.7	36.8	34.7
------	------	------	------	------

% WARE: 3.81 CM (1.5 INCH) RIDDLE

94.1	93.1	95.1	95.3	94.4
------	------	------	------	------

PERMANENT BARLEY BLOCKS

TOTAL TUBERS: TONNES/HECTARE

46.0	42.9	44.3	48.7	45.5
------	------	------	------	------

% WARE: 3.81 CM (1.5 INCH) RIDDLE

93.5	92.9	94.4	93.8	93.6
------	------	------	------	------

71/W/RN/13

WINTER WHEAT

Crop in					N: KG/HA				
1966	1967	1968	1969	1970	63	126	189	252	Mean
GRAIN: TONNES/HECTARE									
(1) and (2)									
W	W	W	L	P	3.22	4.77	4.97	4.63	4.40
W	W	L	P	W	2.70	3.95	3.96	3.59	3.55
W	L	P	W	W	2.09	3.53	4.03	4.19	3.46
W	W	W	W	W	2.88	3.55	3.85	3.17	3.36
Mean (± 0.138)					2.72	3.95	4.20	3.90	3.69

(1) (± 0.317) For use in vertical and diagonal comparisons only
(2) (± 0.276) For use in horizontal and interaction comparisons only

Mean D.M. %: 82.1

STRAW: TONNES/HECTARE									
W	W	W	L	P	3.32	4.06	4.15	3.70	3.81
W	W	L	P	W	2.64	3.87	3.95	3.82	3.57
W	L	P	W	W	2.58	3.38	3.96	4.41	3.58
W	W	W	W	W	3.02	3.66	4.04	3.77	3.62
Mean					2.89	3.74	4.02	3.93	3.64

Mean D.M. %: 87.1

71/W/RN/13

BARLEY

Crop in					N: KG/HA				
1966	1967	1968	1969	1970	63	126	189	252	Mean
GRAIN: TONNES/HECTARE									
(1) and (2)									
B	B	B	L	P	3.59	4.58	4.40	4.82	4.35
B	B	L	P	B	3.82	4.55	5.14	5.36	4.72
B	L	P	B	B	3.71	4.64	4.82	4.69	4.47
B	B	B	B	B	3.07	3.62	4.34	4.64	3.92
Mean (± 0.069)					3.55	4.35	4.67	4.88	4.36

(1) (± 0.189) For use in vertical and diagonal comparisons only

(2) (± 0.138) For use in horizontal and interaction comparisons only

Mean D.M. %: 82.4

STRAW: TONNES/HECTARE									
B	B	B	L	P	2.48	3.30	3.55	3.94	3.32
B	B	L	P	B	2.16	2.87	3.94	3.92	3.22
B	L	P	B	B	2.20	3.17	3.71	3.62	3.18
B	B	B	B	B	1.85	2.83	3.42	3.91	3.00
Mean					2.17	3.04	3.66	3.85	3.18

Mean D.M. %: 87.4