

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1971

[Full Table of Content](#)



---

## 71/R/CS/6 Wheat After Intensive Barley - Wheat

### Rothamsted Research

Rothamsted Research (1972) *71/R/CS/6 Wheat After Intensive Barley - Wheat* ; Yields Of The Field Experiments 1971, pp 130 - 131 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/CS/6

WHEAT AFTER INTENSIVE BARLEY

Object: To study the effects of different periods of pre-cropping with barley on yields and incidence of take-all (*Ophiobolus graminis*) in wheat - Little Knott I.

The eleventh year, winter wheat.

For previous years see 61/C/8(t), 62/C/7, 63-66/C/2, 67/C/2(t), 68/C/2(t), 69/R/CS/6(t) and 70/R/CS/6(t).

Whole plot dimensions: 4.27 x 20.1. Sub plot area harvested: 0.00266.

Blocks were split across all plots for a test of none (U) v. ground chalk at 12.6 tonnes (L).

Basal applications: Manures: 251 kg (0:14:28) combine drilled, 605 kg 'Nitro-Chalk' in spring. Weedkillers: Paraquat at 0.84 kg ion in 225 l. Terbutryne and related triazines ('Prebane' at 4.48 kg in 225 l).

Cultivations, etc.: Paraquat applied: 11 Sept, 1970. Ground chalk applied to half plots: 12 Sept. Ploughed: 23 Sept. Seed drilled at 179 kg: 7 Oct. 'Prebane' applied: 10 Oct. N applied: 13 Apr, 1971. Combine harvested: 25 Aug. Variety: Joss Cambier.

NOTE: Estimates of take-all (*Ophiobolus graminis*) were made in April and early July.

Standard errors per plot.

Grain, tonnes/hectare: Whole plot: 0.461 or 7.7% (39 d.f.)  
Sub plot: 0.626 or 10.4% (40 d.f.)

71/R/CS/6

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

Crop Sequences

Crop in 1961	1	2	3	4	5	6	7	8	9	10	Mean
62	O	WS	O	BE	WS	WS	B	WS	WS	BE	
63	BE	O	WS	O	BE	WS	B	WS	WW	WW	
64	B	BE	O	WS	O	BE	B	WS	WW	P	
65	B	B	BE	O	WS	O	B	WS	WW	B	
66	B	B	B	BE	O	WS	B	WS	WW	BE	
67	B	B	B	B	BE	O	B	WS	WW	WW	
68	B	B	B	B	B	BE	B	WS	F	B	
69	WW	WW	WW	WW	WW	WW	WW	WW	WW	F	
70	F	WW	WW	WW	WW	WW	WW	WW	WW	WW	
RESIDUAL	(±0.326)					(±0.103)					
N3	6.55	6.02	5.77	6.17	6.14	5.56	5.89	5.97	4.80	6.32	5.92
N5	6.91	5.79	6.25	6.34	6.00	6.01	6.16	6.00	4.96	6.10	6.05
N7	7.00	6.27	6.11	6.18	5.61	5.61	6.35	5.95	6.26	6.04	6.14
N9	7.01	6.29	5.35	5.95	5.89	5.91	5.85	5.94	5.21	6.09	5.95
	(±0.221)*										
U	6.72	5.95	5.95	6.20	5.86	5.74	6.01	6.14	5.33	6.48	6.04
L	7.01	6.24	5.78	6.12	5.96	5.81	6.12	5.79	5.29	5.80	5.99
Mean (±0.163)	6.87	6.09	5.87	6.16	5.91	5.77	6.06	5.96	5.31	6.14	6.01

	RESIDUAL			
	N3	N5	N7	N9
	(±0.140)*			
U	5.88	6.19	6.11	5.98
L	5.95	5.92	6.17	5.92

\* For use in horizontal and interaction comparisons only

Mean D.M. %: 81.8