


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/W/CS/56 Nematodes and Verticillium - Potatoes

Rothamsted Research

Rothamsted Research (1972) *71/W/CS/56 Nematodes and Verticillium - Potatoes* ; Yields Of The Field Experiments 1971, pp 190 - 191 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/W/CS/56

NEMATODES AND VERTICILLIUM

Object: To study the effects, direct and residual of methyl bromide, residual only of aldicarb, on *Heterodera rostochiensis* and *Verticillium* on potatoes grown continuously for three years - Woburn Broad Mead I.

The third year, potatoes.

For previous years see 69/W/P/2(t) and 70/W/CS/56(t).

Design: 4 blocks of 6 plots.

Whole plot dimensions: 2.84 x 13.4. Area harvested: 0.00147.

Treatments:

Chemicals 1971: 487 (M1), 975 (M2) kg methyl bromide.

NOTE: These treatments were balanced over (OW) and (BO) applied in 1969. No fresh treatments were applied in 1970.

Basal applications: 1530 kg (13:13:20), 605 kg Epsom salts. Weedkiller: Linuron at 1.12 kg in 281 l. Fungicide with insecticide: Mancozeb at 1.34 kg with demeton-s-methyl at 246 ml in 416 l. Fungicide: Mancozeb at 1.34 kg in 371 l.

Cultivations, etc.: Deep-tine cultivated 1st stroke: 29 Oct, 1970, 2nd stroke: 30 Oct, 3rd stroke: 18 Dec. Methyl bromide applied under gas tight sheets: 21 Apr, 1971. Gas tight sheets removed: 26 Apr. NPK applied: 27 Apr. Mg applied, rotary cultivated, potatoes planted: 28 Apr. Weedkiller applied: 17 May. Rotary ridged twice: 21 June, 25 June. Fungicide with insecticide applied: 29 June. Fungicide applied: 12 Aug. Lifted: 13 Sept. Variety: Pentland Crown.

NOTE: Soil samples were taken after treatments were applied and after harvest for numbers of cysts, eggs and larvae of *Heterodera rostochiensis* and propagules of *Verticillium*.

Standard error per plot.

Total tubers, tonnes/hectare: 3.75 or 21.6% (15 d.f.)

71/W/CS/56

SUMMARY OF RESULTS

00	1969				1971		Mean
	MD	MW	TO		M1	M2	
TOTAL TUBERS: TONNES/HECTARE							
(±1.88)							
5.9	6.2	7.7	8.8	37.3	38.1	17.3	
% WARE: 4.44 CM (1.75 INCH) RIDDLE							
29.7	33.9	45.2	42.5	87.5	89.8	54.8	