

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/CS/59 Break Crops and Wheat - Wheat

Rothamsted Research

Rothamsted Research (1972) *71/R/CS/59 Break Crops and Wheat - Wheat* ; Yields Of The Field Experiments 1971, pp 195 - 196 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/CS/59

BREAK CROPS AND WHEAT

Object: To study the effects of different break crops on yields and soil-borne pathogens of a following sequence of cereals - Geescroft.

The second year, winter wheat.

For previous year see 70/R/CS/59(t).

Design: 3 randomised blocks of 14 plots, split into 4.

Whole plot dimensions: 4.27 x 33.2. Sub plot area harvested: 0.00215.

Treatments:-

Whole plots: Crops and nitrogen 1970:-

Maize receiving N at 100 (MA2), 200 (MA4) kg as 'Nitro-Chalk' in seedbed, spring beans (2 plots per block) (BE), clover (2 plots per block) (CL).
and all combinations of:-

1. Crops: Barley (B), oats (O).
2. Undersowing: None (O), trefoil (T).
3. Nitrogen: 50 (1), 100 (2) kg as 'Nitro-Chalk' in seedbed.

Sub plots:

4. Nitrogen 1971: None, 50, 100, 150 kg as 'Nitro-Chalk'.

Basal and other applications: Corrective K to clover plots at 126 kg K₂O as muriate of potash. 314 kg (0:20:20) combine drilled.
Weedkiller: 2,4-D at 0.56 kg and dichlorprop at 2.24 kg in 225 l.

Cultivations, etc.: Corrective K applied: 20 Oct, 1970. Ploughed: 3 Nov. Seed combine drilled at 202 kg: 5 Nov. N applied: 8 Apr, 1971. Weedkiller applied: 14 Apr. Combine harvested: 25 Aug. Variety: Joss Cambier.

NOTE: Estimates were made of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*).

Standard errors per plot.

Grain, tonnes/hectare: Whole plot: 0.238 or 4.9% (26 d.f.)
Sub plot: 0.363 or 7.5% (84 d.f.)

7L/R/CS/59

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

N: KG/HA 1971

		0	50	100	150	Mean
Crop and N 1970	Under-sowing 1970	(1) and (2)				(±0.137)
B1	O	2.73	3.94	4.32	4.26	3.81
B2	O	2.51	3.90	4.13	4.61	3.79
B1	T	3.38	3.74	4.48	4.20	3.95
B2	T	3.73	4.47	4.36	4.42	4.25
O1	O	3.52	4.97	5.50	5.36	4.84
O2	O	4.06	5.66	5.71	5.42	5.21
O1	T	3.65	5.27	5.66	5.18	4.94
O2	T	4.40	5.10	5.60	5.36	5.11
MA2	O	4.79	5.59	5.72	5.07	5.29
MA4	O	5.20	5.70	5.63	5.10	5.41
		(3) and (4)				(±0.097)
BE*	O	4.53	5.72	5.81	5.54	5.40
CL*	O	4.49	5.56	5.50	5.16	5.18
Mean (±0.056)		4.00	5.07	5.27	5.03	4.84

Mean D.M. %: 83.9

(1) (±0.228) (3) (±0.161) For use in vertical and diagonal comparisons only

(2) (±0.210) (4) (±0.148) For use in horizontal and interaction comparisons only

* Duplicated treatments