Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1971



Full Table of Content

71/W/CS/63 Nematodes and Verticillium - Potatoes

Rothamsted Research

Rothamsted Research (1972) 71/W/CS/63 Nematodes and Verticillium - Potatoes; Yields Of The Field Experiments 1971, pp 202 - 203 - DOI: https://doi.org/10.23637/ERADOC-1-97

71/W/CS/63

NEMATODES AND VERTICILLIUM

Object: To study the residual effects of methyl bromide, aldicarb and benomyl (applied to potatoes in 1970) on Heterodera rostochiensis and Verticillium on a second potato crop in 1971 - Woburn Broadmead I.

The second year, potatoes.

For previous year see 70/W/P/2(t).

Design: 4 blocks of 4 plots.

Whole plot dimensions: 2.84 x 13.4. Area harvested: 0.00147.

Treatments:

Fresh treatments were not applied in 1971.

- Basal applications: 1530 kg (13:13:20). 897 kg Epsom salts. Weedkillers: Linuron at 1.12 kg in 281 l. Paraquat was applied at a concentration of 0.03 kg ion in 4.55 l to the inter-row area rate per hectare not recorded. Fungicide with insecticide: Mancozeb at 1.34 kg with demeton-s-methyl at 246 ml in 416 l.
- Cultivations, etc.: Deep-tine cultivated 1st stroke: 27 Oct, 1970, 2nd stroke: 30 Oct, 3rd stroke: 18 Dec. NPK and Mg applied, rotary cultivated, potatoes planted: 27 Apr, 1971. Linuron applied: 17 May. Paraquat applied to inter-row area: 7 8 June. Rotary ridged 1st stroke: 21 June, 2nd stroke: 25 June. Fungicide and insecticide applied: 29 June. Lifted: 13 Sept. Variety: Pentland Dell.
- NOTE: Soil samples were taken before planting and after harvest for number of cysts, eggs and larvae of Heterodera rostochiensis and propagules of Verticillium.
- Standard error per plot.
 Total tubers, tonnes/hectare: 3.01 or 25.4% (9 d.f.)

71/W/CS/63 SUMMARY OF RESULTS

0	В	М	T	Mean	
	TO	TAL TUBERS: TO	NNES/HECTARE		-
	(±1.51)				
3.5	18.4	15.2	10.3	11.9	
	% WAR	E: 4.44 CM (1.	75 INCH) RIDDI	E	
0.7	56.9	44.5	30.5	33.1	