Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1971



Full Table of Content

71/W/CS/78 Nematodes and Verticillium - Potatoes

Rothamsted Research

Rothamsted Research (1972) 71/W/CS/78 Nematodes and Verticillium - Potatoes; Yields Of The Field Experiments 1971, pp 229 - 230 - DOI: https://doi.org/10.23637/ERADOC-1-97

71/W/CS/78

NEMATODES AND VERTICILLIUM

Object: To study the effects of methyl bromide, aldicarb, benomyl and dazomet on Heterodera rostochiensis and Verticillium on potatoes - Woburn Broadmead I.

The first year, potatoes.

Design: 4 blocks of 6 plots.

Whole plot dimensions: 2.84 x 12.8. Area harvested: 0.00147.

Treatments:

Chemicals: None (0), aldicarb at 6.7 kg (A), benomyl at 22.4 kg (B), aldicarb at 6.7 kg plus benomyl at 22.4 kg (A+B), dazomet at 336 kg (D), methyl bromide at 975 kg (M).

Basal applications: 1534 kg (13:13:20), 605 kg Epsom salts. Weedkiller: Linuron at 1.12 kg in 281 l. Fungicide with insecticide: Mancozeb at 1.34 kg with demeton-s-methyl at 246 ml in 416 l. Fungicide: Mancozeb at 1.34 kg in 371 l.

Cultivations, etc.: Deep-tine cultivated 1st stroke: 27 Oct, 1970, 2nd stroke: 30 Oct, 3rd stroke: 18 Dec. Dazomet applied, all plots rotary cultivated: 23 Mar, 1971. Methyl bromide applied under gas tight sheets: 22 Apr. Gas tight sheets removed: 26 Apr. Aldicarb and benomyl applied, NPK and Mg applied, rotary cultivated, potatoes planted: 27 Apr. Weedkiller applied: 17 May. Rotary ridged: 21 June. Fungicide with insecticide applied: 29 June. Fungicide applied: 13 Aug. Lifted: 13 Sept. Variety: Pentland Dell. Previous crops: Barley 1969 and potatoes 1970.

NOTE: Soil samples were taken after treatments were applied and after harvest for numbers of cysts eggs and larvae of Heterodera rostochiensis and propagules of Verticillium.

Standard error per plot.
Total tubers, tonnes/hectare: 3.09 or 9.0% (15 d.f.)

71/W/CS/78 SUMMARY OF RESULTS

0	A	В	A+B	D	M	Mean
		TOTAL T	UBERS: TO	NNES/HECTA	RE	-
		(±1.	55)			
21.5	33.8	35.1	36.0	36.3	43.9	34.4
		% WARE: 4	.바 CM (1.	75 INCH)	RIDDLE	
65.8	78.8	82.5	81.1	77.6	82.4	78.0