

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/B/1 and 71/BB/B/1 Growth and Yield on Contrasted Sites - Barley

Rothamsted Research

Rothamsted Research (1972) *71/R/B/1 and 71/BB/B/1 Growth and Yield on Contrasted Sites - Barley* ; Yields Of The Field Experiments 1971, pp 263 - 266 - DOI:

<https://doi.org/10.23637/ERADOC-1-97>

71/R/B/1 and 71/BB/B/1

BARLEY

GROWTH AND YIELD ON CONTRASTED SITES

Object: To try to account for yields and differences between yields of barley on sites at Rothamsted and Broom's Barn by studying crop growth rates, nutrient uptake, water use etc. at a wide range of nitrogen levels, with and without irrigation. Also to study the interaction between site differences and crops (see also 71/R&BB/WW/6)- Rothamsted (R) Great Knott III and Broom's Barn (BB) Marl Pit Field.

Design: 3 randomised blocks of 2 plots, split into 6.

Whole plot dimensions:-

Great Knott III (R): 15.2 x 48.0. Sub plot area harvested: 0.00434.
Marl Pit Field (BB): 15.2 x 45.7. Sub plot area harvested: 0.00413.

Treatments: All combinations of:-

Whole plots: 1. Irrigation: None (0), full irrigation (I).
Sub plots: 2. Nitrogen: 31, 63, 94, 125, 157, 188 kg N as 'Nitro-Chalk'.

Total irrigation was 33.0 mm applied on two occasions (R) and 58.4 mm applied on three occasions (BB).

Basal applications:

Great Knott III (R): 1260 kg (0:20:20) plus 900 kg Epsom salts ploughed in in autumn, 360 kg (0:20:20) combine drilled.
Marl Pit Field (BB): 1260 kg (0:20:20) plus 900 kg Epsom salts ploughed in in autumn 251 kg (0:20:20) combine drilled.
Weedkiller (both fields): MCPA, mecoprop and dicamba ('Banlene Plus' at 5.6 l in 225 l).
Fungicide (both fields): Tridemorph at 0.53 kg applied with the weedkiller.
Weedkiller to Great Knott III (R) only: Paraquat at 0.56 kg ion in 225 l.

Cultivations, etc.:

Great Knott III (R): Basal PK and Epsom salts applied: 15 Sept, 1970. Paraquat applied: 14 Oct. Ploughed: 2 - 10 Nov. N applied, seed combine drilled at 180 kg: 11 Mar, 1971. Weedkiller plus fungicide applied: 10 May. Irrigated: 20 May - 5.1 mm, 8 June - 27.9 mm. Combine harvested: 18 Aug. Variety: Julia. Previous crops: Potatoes 1969, winter wheat 1970.

71/R/B/1 and 71/BB/B/1

Marl Pit Field (BB): Basal PK applied: 18 Sept, 1970. Epsom salts applied: 16 Oct. Ploughed: 20 Oct. Seed combine drilled at 180 kg: 5 Mar, 1971. N applied: 23 Mar. Weedkiller plus fungicide applied: 10 May. Irrigated: 14 May - 26.7 mm, 21 May - 16.5 mm, 26 May - 15.2 mm. Combine harvested: 12 Aug. Variety: Julia. Previous crops: Sugar beet 1969, barley 1970.

Standard errors per plot. Grain, tonnes/hectare:

Great Knott III (R): Sub plot: 0.303 or 4.8% (20 d.f.)

Marl Pit Field (BB): Sub plot: 0.266 or 4.9% (20 d.f.)

71/R/B/1 and 71/BB/B/1

SUMMARY OF RESULTS

GREAT KNOTT III (R)

N: KG/HA

	31	63	94	125	157	188	Mean
GRAIN: TONNES/HECTARE							
(±0.175)*							
O	5.65	6.40	6.71	6.64	6.64	6.21	6.37
I	5.36	6.42	6.56	6.60	6.85	6.19	6.33
Mean (±0.124)	5.50	6.41	6.64	6.62	6.75	6.20	6.35
STRAW: TONNES/HECTARE							
O	3.25	5.35	5.60	5.77	5.90	5.58	5.24
I	3.61	5.11	5.89	5.65	6.34	5.88	5.41
Mean	3.43	5.23	5.74	5.71	6.12	5.73	5.33

Mean D.M. %: Grain: 83.5
 Straw: 77.6

* For use in horizontal and interaction comparisons only

71/R/B/1 and 71/BB/B/1

MARL PIT FIELD (BB)

N: KG/HA

	31	63	94	125	157	188	Mean
GRAIN: TONNES/HECTARE							
(±0.154)*							
O	4.07	4.99	5.37	5.85	5.90	5.77	5.33
I	4.76	5.34	6.01	5.93	6.11	5.56	5.62
Mean (±0.109)	4.41	5.16	5.69	5.89	6.00	5.67	5.47
STRAW: TONNES/HECTARE							
O	3.30	4.28	4.81	5.56	5.72	6.08	4.96
I	3.97	5.04	6.08	6.25	6.72	6.36	5.74
Mean	3.63	4.66	5.45	5.91	6.22	6.22	5.35

Mean D.M. %: Grain: 82.7
 Straw: 82.8

* For use in horizontal and interaction comparisons only