

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/B/10 Systemic Fungicides - Barley

Rothamsted Research

Rothamsted Research (1972) *71/R/B/10 Systemic Fungicides - Barley* ; Yields Of The Field Experiments 1971, pp 285 - 286 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/B/10

BARLEY

SYSTEMIC FUNGICIDES

Object: To study the effects of a range of fungicides on yield and pathogens of barley - Drapers.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 4.27 x 13.4. Area harvested: 0.00190.

Treatments:

No fungicidal treatment	(O)
Seed dressings per kg of seed:	
3191 F at 2 g	(BA)
3191 F plus confidential ingredient, at 3 g	(BAC)
3200 F at 4 g	(BB)
Organo-mercury at 2.2 g	(ME)
MC 833 at 2.0 g	(MC)
'Vitavax' plus thiram, at 2.2 g	(VT)
NF 48 at 24 g	(NF)
'Vitavax' plus organo-mercury, at 2.2 g	(V)
Soil drench after sowing:	
Sandoz Fungicide at 11.0 l a.i. in 5600 l	(SF)

Basal applications: 505 kg (20:10:10) combine drilled. Weedkillers:
Paraquat at 0.28 kg ion in 225 l.

Cultivations, etc.: Paraquat applied: 23 Sept, 1970. Ploughed: 3 Oct.
Seed combine drilled at 157 kg: 26 Mar, 1971. Soil drench applied:
29 Mar. Combine harvested: 19 Aug. Variety: Sultan. Previous
crops: Barley 1969 and 1970.

NOTE: Samples were taken for mildew (*Erysiphe graminis*), brown rust
(*Puccinia hordei*), smut (*Ustilago nuda*) and root diseases.

Standard error per plot.

Grain, tonnes/hectare: 0.285 or 7.3% (18 d.f.)

71/R/B/10

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

O	BA	BAC	BB	ME	MC	VT	NF	V	SF	Mean
										(±0.165)
3.69	3.66	3.09	4.38	3.95	4.01	3.63	4.83	4.11	3.72	3.91

Mean D.M.%: 82.0