

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1971

[Full Table of Content](#)



### 71/R/B/13 Methods of Applying N P K - Barley

#### Rothamsted Research

Rothamsted Research (1972) *71/R/B/13 Methods of Applying N P K - Barley* ; Yields Of The Field Experiments 1971, pp 291 - 292 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/B/13

BARLEY

METHODS OF APPLYING NPK

Object: To study the effect of injecting NPK fertiliser between bands of seed, using the Tume Combi drill, on the yields of barley - West Barnfield I.

Design: 6 randomised blocks of 4 plots.

Dimensions of whole plot: 4.88 x 15.2. Area harvested: 0.00464.

Treatments: All combinations of:-

1. Drills and row spacing: Combine drill, 17.8 cm (7 inches) between rows (C), Tume Combi drill 25.4 cm (10 inches) between band centres (T).
2. Rates of compound fertiliser (25:10:10): 264, 505 kg.

Basal applications: No manures. Weedkillers: Paraquat at 0.28 kg ion in 225 l and later at 0.56 kg ion in 225 l, ioxynil octanoate, bromoxynil octanoate and the iso-octyl ester of dichlorprop ('Oxytril P' at 1.4 l in 225 l).

Cultivations, etc.: Paraquat applied: 19 Sept, 1970. Ploughed: 24 Sept. Paraquat applied: 5 Feb, 1971. Seed combine drilled at 157 kg: 26 Mar. 'Oxytril P' applied: 11 May. Combine harvested: 17 Aug. Variety: Julia. Previous crops: Spring oilseed rape 1969, winter wheat 1970.

Standard error per plot.

Grain, tonnes/hectare: 0.276 or 5.4% (15 d.f.)

71/R/B/13

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

(25:10:10): KG/HA

	264	505	Mean
	( $\pm 0.113$ )		( $\pm 0.080$ )
C	5.03	5.14	5.09
T	4.99	5.17	5.08
Mean ( $\pm 0.080$ )	5.01	5.16	5.09

Mean D.M. %: 85.0