


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/S/B/1 Varieties, N Rates and Times of Application - Barley

Rothamsted Research

Rothamsted Research (1972) *71/S/B/1 Varieties, N Rates and Times of Application - Barley* ; Yields Of The Field Experiments 1971, pp 293 - 295 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/s/B/1

BARLEY

VARIETIES, N RATES AND TIMES OF APPLICATION

Object: To study the effects of a range of nitrogen levels applied to the seedbed or in May, on the yield of two barley varieties. The effects of dressing seed with phenylphosphonic acid are also studied - Saxmundham Grove Plot.

Design: Two randomised blocks of 6 plots split into 4.

Whole plot dimensions: 2.74 x 12.2. Sub plot area harvested: 0.00052.

Treatments:-

Whole plots: All combinations of:-

1. Varieties: Midas (M), Sultan (S).
2. N levels: 50, 100, 150 kg N as calcium nitrate.

Sub plots: All combinations of:-

3. Times of applying N: To seedbed (E), top-dressed in May (L), (across whole plots).
4. Seed dressing: None (O), phenylphosphonic acid (0.2 % a.i.) sprayed over the seed as a seed dressing (D) (along whole plots).

Basal applications: 314 kg (0:20:20) broadcast. Weedkiller: Dichlorprop plus MCPA ('Mephetol Plus' at 5.61 l in 562 l). Fungicide: Tridemorph at 0.53 kg in 562 l.

Cultivations, etc.: Ploughed: 28 Oct, 1970. Seed drilled at 188 kg, seedbed N and basal PK applied: 31 Mar, 1971. Weedkiller applied: 18 May. Late N applied: 19 May. Fungicide applied: 17 June. Harvested by sickle: 11 Aug. Previous crops: Barley 1969, sugar beet 1970.

Standard errors per plot.

Grain, tonnes/hectare: Whole plot: 0.164 or 6.4% (5 d.f.)

Sub plot (T): 0.410 or 16.3% (6 d.f.)

(S): 0.231 or 9.2% (6 d.f.)

(TS): 0.289 or 11.5% (6 d.f.)

Pooled used for calculation of standard error in the summary:

0.320 or 12.7% (23 d.f.)

71/S/B/1

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

	N: KG/HA			E	L	O	D	Mean
	50	100	150					
	(±0.113)			(±0.092)		(±0.092)		(±0.065)
M	2.47	2.50	2.37	2.62	2.28	2.32	2.58	2.45
S	2.64	2.68	2.45	2.70	2.48	2.52	2.66	2.59
	N: KG/HA			(±0.113)		(±0.113)		(±0.080)
			50	2.65	2.45	2.44	2.66	2.55
			100	2.79	2.39	2.44	2.74	2.59
			150	2.53	2.29	2.37	2.46	2.41
						(±0.092)		(±0.065)
					E	2.50	2.82	2.66
					L	2.33	2.43	2.38
Mean (±0.047)						2.42	2.62	2.52

Mean D.M. %: 76.4

71/S/B/1

STRAW: TONNES/HECTARE

	N: KG/HA			E	L	O	D	Mean
	50	100	150					
M	3.88	4.56	4.54	4.55	4.11	4.22	4.43	4.33
S	3.52	3.92	4.23	3.96	3.82	3.69	4.09	3.89
		N: KG/HA						
			50	3.85	3.55	3.70	3.71	3.70
			100	4.55	3.93	4.09	4.39	4.24
			150	4.36	4.41	4.09	4.69	4.39
					E	4.09	4.42	4.25
					L	3.83	4.10	3.96
Mean						3.96	4.26	4.11

Mean D.M. %: 69.9