

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1971

[Full Table of Content](#)



---

### 71/R/BE/3 Control of Vectors and Viruses - Beans

#### Rothamsted Research

Rothamsted Research (1972) *71/R/BE/3 Control of Vectors and Viruses - Beans* ; Yields Of The Field Experiments 1971, pp 304 - 305 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/BE/3

SPRING BEANS

CONTROL OF VECTORS AND VIRUSES

Object: To study the effects of a range of insecticide treatments on yield and incidence of weevils, aphids and virus diseases of beans - Long Hoos VI.

Design: 4 randomised blocks of 6 plots.

Whole plot dimension: 10.7 x 12.2. Area harvested: 0.00293.

Treatments: Insecticides:

None	(O)
Applied to soil:-	
Aldicarb 1.12 kg	(AS1)
Aldicarb 11.2 kg	(AS2)
Foliar sprays on 3 occasions:-	
BHC 0.56 kg	(BF)
Malathion 1.12 kg	(MAF)
Menazon 0.28 kg	(MEF)

The first and second sprays were applied in 524 l, the third in 700 l.

Basal applications: 7530 kg ground chalk. 405 kg (0:14:28) placement drilled. Weedkiller: Simazine at 1.12 kg in 225 l.

Cultivations, etc.: Ground chalk applied: 21 Sept, 1970. Ploughed: 30 Oct. Aldicarb applied, all plots rotary cultivated, seed placement drilled at 224 kg: 1 Mar, 1971. Weedkiller applied: 11 Mar. Foliar sprays applied: 27 Apr, 18 May, 30 June. Combine harvested: 3 Sept. Variety: Maris Bead. Previous crops: Lucerne and Cocksfoot (fallowed after 1st cut) 1969, winter wheat 1970.

NOTE: Aphid and weevil counts were made and virus incidence assessed.

Standard error per plot.

Grain, tonnes/hectare: 0.232 or 7.6% (15 d.f.)

71/R/BE/3

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

0	AS1	AS2	BF	MAF	MEF	Mean
		(±0.116)				
2.84	3.04	3.30	2.96	3.14	3.12	3.07

Mean D.M. %: 80.7