

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1971

[Full Table of Content](#)



## 71/R/SC/1 Seed Spacing and N - Sweet Corn

### Rothamsted Research

Rothamsted Research (1972) *71/R/SC/1 Seed Spacing and N - Sweet Corn* ; Yields Of The Field Experiments 1971, pp 335 - 335 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/SC/1

SWEET CORN

SEED SPACING AND N

Object: To study the effects of seed spacing and levels of nitrogen on yield of sweet corn - Long Hoos V 1.

Design: 4 randomised blocks of 12 plots.

Whole plot dimensions: 2.44 x 5.18.

Treatments: All combinations of:

1. Seed spacing: Seed 7.6 cm (3 inches), 15.2 cm (6 inches) 30.5 cm (12 inches) apart in rows 61.0 cm (2 feet) apart.
2. Nitrogen: None, 63, 126, 188 kg N as 'Nitro-Chalk'.

Basal applications: 1255 kg (0:14:28). Weedkiller: Atrazine at 1.68 kg in 337 l. Insecticide: Phorate granules at 1.8 kg a.i. applied by hand-blower.

Cultivations, etc.: Basal PK applied: 27 Oct, 1970. Ploughed: 29 Oct. Rotary cultivated: 4 May, 1971. Seed precision drilled: 5 May. N applied: 18 May. Weedkiller and insecticide applied: 21 May. Harvested: 16-28 Sept. Variety: Early King. Previous crops: Potatoes 1969, barley 1970.

NOTE: Because phorate granules were irregularly applied a subsequent attack of frit fly (*Oscinella frit*) was only controlled on parts of plots. Yields were therefore not taken.