

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1971

[Full Table of Content](#)



### 71/R/K/1 Virus Control - Kale

#### Rothamsted Research

Rothamsted Research (1972) *71/R/K/1 Virus Control - Kale* ; Yields Of The Field Experiments 1971, pp 336 - 337 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/K/1

KALE

VIRUS CONTROL

Object: To study the effects of insecticides on virus infection and yield of kale - Great Field I.

Design: 4 randomised blocks of 4 plots.

Whole plot dimensions: 10.7 x 10.7. Area harvested: 0.00114.

Treatments: Insecticides:-

All combinations of:

(1) Menazon: None (0), 'Saphicol' at 1.05 l in 674 l (M)

(2) BHC: None (0), 0.21 kg in 674 l (B)

The treatments were applied on 3 occasions (17 June, 16 July, 19 Aug).

Basal applications: 251 kg (0:20:20), 942 kg (20:10:10). Weedkiller: Linuron at 0.84 kg in 427 l.

Cultivations, etc.: Ploughed: 2 Nov, 1970. PK applied: 3 Apr, 1971. NPK applied: 5 Apr. Seed drilled at 2.24 kg: 7 Apr. Weedkiller applied: 22 May. Singled: 2 June. Harvested: 17 Dec. Variety: Maris Kestrel. Previous crops: Spring wheat 1969, spring beans 1970.

NOTE: Observations were made on aphid infestation and virus infection.

Standard error per plot.

Fresh weight, tonnes/hectare: 10.01 or 13.4% (9 d.f.)

71/R/K/1

SUMMARY OF RESULTS

FRESH WEIGHT: TONNES/HECTARE

	O	B	Mean
	(±5.01)		(±3.54)
O	76.2	76.1	76.1
M	72.5	74.0	73.3
Mean (±3.54)	74.3	75.0	74.7