

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1971

[Full Table of Content](#)



### 71/R/M/5 Amidophosphates for Barley (Resown With Ryegrass), Potatoes and Kale - Mixed Crops

#### Rothamsted Research

Rothamsted Research (1972) *71/R/M/5 Amidophosphates for Barley (Resown With Ryegrass), Potatoes and Kale - Mixed Crops* ; Yields Of The Field Experiments 1971, pp 348 - 351 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

TL/R/M/5

MIXED CROPS

AMIDOPHOSPHATES FOR BARLEY (RESOWN WITH RYEGRASS), POTATOES AND KALE

Object: To compare different forms of NP fertiliser on a range of crops - West Barnfield I.

Design: 3 randomised blocks of 9 plots for each crop.

Whole plot dimensions:

Barley (resown with ryegrass): 3.66 x 1.83. Area harvested:  
1st cut, 0.00037, 2nd cut, 0.00038, 3rd cut, 0.00028.  
Potatoes: 3.66 x 2.85. Area harvested: 0.00048.  
Kale: 3.66 x 2.13. Area harvested: 1st and 2nd cut, 0.00005,  
3rd cut, 0.00026.

Treatments: None (O) and additional treatments DS, TS (Not applied to barley or ryegrass - see note at end),

where T = Phosphoryltri-*am*ide

D = Sodium di-amidophosphate

S = Sprayed on to the crop. Rates: To give 11.2 kg P for potatoes, 8.41 kg P for kale.

and all combinations of:-

1. NP fertilisers: Ammonium phosphate (A), sodium di-amidophosphate (D), urea phosphate (U).

2. Rates of NP fertilisers: To give:-

To barley (resown with ryegrass): 11.2 kg (1), 22.4 kg (2) P.  
To potatoes: 22.4 kg (1), 44.8 kg (2) P.  
To kale: 16.8 kg (1), 33.6 kg (2) P.

Basal applications: Weedkiller: Paraquat at 0.28 kg in 225 l, and later at 0.56 kg ion in 225 l.

Standard applications:

Barley (resown with ryegrass): 56 kg N, 56 kg K in seedbed for barley, 56 kg N to ryegrass after the first cut and 112 kg N after the second.

Potatoes: 112 kg N, 168 kg K. Weedkiller: Linuron at 0.84 kg plus paraquat at 0.42 kg ion in 562 l. Fungicide plus insecticide: Mancozeb at 1.35 kg plus menazon ('Saphicol' at 0.70 l) in 449 l on 3 occasions.

Kale: 112 kg N, 112 kg K, plus 112 kg N in mid-season.

Insecticide: Menazon ('Saphicol' at 0.70 l in 449 l).

For all crops N was applied as 'Nitro-Chalk', K as muriate of potash.

71/R/M/5

2

Cultivations, etc.: Paraquat applied: 19 Sept, 1970. Ploughed: 24 Sept. Paraquat applied: 5 Feb, 1971.  
Barley (resown with ryegrass): NP and NK fertilisers applied, barley seed drilled at 1.91 kg: 1 Apr, 1971. Resown with ryegrass: 28 Apr. Cut, N applied: 8 June. Cut: 14 July. N applied: 23 July. Cut: 7 Sept. Variety: Western Wolths. (Barley: Julia).  
Potatoes: NP and NK fertilisers applied, potatoes planted: 13 Apr, 1971. Weedkiller applied: 12 May. Fungicide plus insecticide applied: 16 June. Amidophosphate sprays applied: 22 June. Fungicide plus insecticide applied: 6 July, 28 July. Lifted: 1 Sept. Variety: King Edward.  
Kale: NP and NK fertilisers applied, seed drilled: 2 Apr, 1971. Amidophosphate sprays applied, midseason N applied: 22 June. Insecticide applied: 6 July. Harvested: 5 Oct. Variety: Thousand Headed.  
Previous crops: Spring Oilseed Rape 1969, winter wheat, 1970.

NOTE: After sowing the barley there was considerable bird damage. Ryegrass was sown between the rows and the barley cut as a green crop. Amidophosphate sprays were not applied to these crops.

Standard errors per plot.

Barley (resown with ryegrass): Grain, tonnes/hectare:  
1st cut: 0.247 or 21.9% (16 d.f.)  
2nd cut: 0.240 or 7.6% (16 d.f.)  
3rd cut: 0.123 or 6.4% (16 d.f.)  
Total of 3 cuts: 0.266 or 4.3% (16 d.f.)  
Potatoes: Total tubers, tonnes/hectare: 3.20 or 8.9% (16 d.f.)  
Kale: Dry matter, tonnes/hectare: 0.942 or 9.8% (16 d.f.)

71/R/M/5

SUMMARY OF RESULTS

BARLEY AND RYEGRASS, DRY MATTER: TONNES/HECTARE

D	A1	A2	D1	D2	U1	U2	Mean
1ST CUT							
0.91 (±0.082)	1.18	1.19	1.42 (±0.143)	1.55	0.99	1.07	1.13
2ND CUT							
3.21 (±0.080)	2.96	3.29	2.97 (±0.139)	3.24	3.06	3.20	3.15
3RD CUT							
1.85 (±0.041)	2.03	1.90	1.84 (±0.071)	2.05	1.93	2.02	1.92
TOTAL OF 3 CUTS							
5.98 (±0.089)	6.17	6.37	6.22 (±0.153)	6.84	5.98	6.30	6.20

Mean D.M. %: 1st cut: 15.5  
 2nd cut: 22.4  
 3rd cut: 37.6  
 Total of 3 cuts: 25.2

71/R/M/5

O	A1	A2	D1	D2	DS	U1	U2	TS	Mean
POTATOES, TOTAL TUBERS: TONNES/HECTARE									
(±1.85)									
34.5	36.2	37.6	37.0	37.2	34.1	31.5	38.7	35.7	35.8

KALE, DRY MATTER: TONNES/HECTARE									
(±0.544)									
8.20	8.97	9.28	9.53	9.92	9.47	9.53	11.56	10.19	9.63