

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/RN/1 and 71/R/RN/2 Ley/ARABLE - Old Grass, Leys, Barley, Wheat

Rothamsted Research

Rothamsted Research (1972) *71/R/RN/1 and 71/R/RN/2 Ley/ARABLE - Old Grass, Leys, Barley, Wheat* ; Yields Of The Field Experiments 1971, pp 50 - 67 - DOI:

<https://doi.org/10.23637/ERADOC-1-97>

71/R/RN/1 and 71/R/RN/2

LEY/ARABLE

Object: To study the effects of three-year leys on the fertility of the soil as measured by a sequence of three arable test crops. Since 1968, continuous winter wheat has been grown after the three test crops to study the build-up and decline of take-all (*Ophiobolus graminis*) after the different cropping sequences - Highfield and Fosters.

The 23rd year, old grass, leys, barley, wheat.

For previous years see 'Details' 1967, 68/B/1(t), 69/R/RN/1&2(t) and 70/R/RN/1&2(t).

Continuous winter wheat: The basal PK for blocks in continuous winter wheat is now 75 kg P205, 75 kg K20 as (0:20:20) combine drilled, the N test is unchanged.

Management of hay plots (H): These plots now receive 75 kg N, 75 kg P205, 75 kg K20 as (15:15:15) in spring and 75 kg N, 48 kg K20 as (25:0:16) after each cut except the last. They are cut at the same times as the leys.

HIGHFIELD

2nd year Treatment Crops:

All-grass ley: PK applied: 20 Oct, 1970. NK applied: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. NK applied after first three cuts.

Clover-grass ley: PK applied: 20 Oct, 1970. K applied: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. K applied after first three cuts.

Lucerne: PK applied: 20 Oct, 1970. Sprayed with paraquat at 0.56 kg ion in 225 l: 4 Feb, 1971. Cut three times: 4 June, 14 July, 7 Sept. Variety: Du Puits.

Hay: Seed undersown in barley at 32 kg: 7 May, 1970. NPK applied: 17 Mar, 1971. Cut four times: 26 May, 14 July, 13 Sept, 2 Nov. NK applied after first three cuts.

2nd Test Crop. Wheat:-

PK applied: 26 Sept, 1970. Deep-tine cultivated twice: 28 Sept. Seed combine drilled at 202 kg: 5 Oct. N applied, plots sprayed with 2,4-D at 0.56 kg plus dichlorprop at 2.24 kg in 225 l: 14 Apr, 1971. Combine harvested: 24 Aug. Variety: Joss Cambier.

71/R/RN/1 and 71/R/RN/2

3rd Test Crop. Barley:-

Ground chalk applied: 11 Sept, 1970. Ploughed twice: 11 Sept, 15 Dec. Seed combine drilled at 157 kg: 25 Feb, 1971. N applied: 5 Mar. Sprayed with ioxynil at 0.53 kg plus mecoprop at 1.57 kg in 225 l: 3 May. Combine harvested: 16 Aug. Variety: Julia.

4th, 6th and 7th Test Crops. Wheat:-

Ploughed: 11 Sept, 1970. Seed combine drilled at 202 kg: 5 Oct. N applied: 13 Apr, 1971. Sprayed with 2,4-D at 0.56 kg plus dichlorprop at 2.24 kg in 225 l: 14 Apr. Combine harvested: 24 Aug. Variety: Joss Cambier.

Permanent Grasses:

The 23rd experimental year permanent (old) grass, blocks 1, 2 and 4, the 23rd year reseeded grass, blocks 1 and 4. PK applied: 20 Oct, 1970. NK applied to 'all-grass' half plots, K to 'clover-grass' half plots: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. NK applied to 'all-grass' half plots and K to 'clover-grass' half plots after each cut except the last.

FOSTERS

2nd year Treatment Crops:

All-grass ley: PK applied: 20 Oct, 1970. NK applied: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. NK applied after first three cuts.

Clover-grass ley: PK applied: 20 Oct, 1970. K applied: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. K applied after first three cuts.

Lucerne: PK applied: 20 Oct, 1970. Sprayed with paraquat at 0.56 kg ion in 225 l: 4 Feb, 1971. Cut three times: 4 June, 14 July, 7 Sept. Variety: Du Puits.

Hay: Seed undersown in barley at 32 kg: 7 May, 1970. NPK applied: 17 Mar, 1971. Cut four times: 26 May, 14 July, 13 Sept, 2 Nov. NK applied after first three cuts.

2nd Test Crop. Wheat:-

PK applied: 26 Sept, 1970. Deep-tine cultivated twice: 28 Sept. Seed combine drilled at 202 kg: 5 Oct. N applied: 13 Apr, 1971. Sprayed with 2,4-D at 0.56 kg plus dichlorprop at 2.24 kg ion 225 l: 15 Apr. Combine harvested: 24 Aug. Variety: Joss Cambier.

71/R/RN/1 and 71/R/RN/2

3rd Test Crop. Barley:-

Ploughed twice: 10 Sept, 14 Dec, 1970. Seed combine drilled at 157 kg: 25 Feb, 1971. N applied: 5 Mar. Sprayed with ioxynil at 0.53 kg plus mecoprop at 1.57 kg in 225 l: 3 May. Combine harvested: 16 Aug. Variety: Julia.

4th, 6th and 7th Test Crops. Wheat:-

Ploughed: 10 Sept, 1970. Seed combine drilled at 202 kg: 6 Oct. N applied: 13 Apr, 1971. Sprayed with 2,4-D at 0.56 kg plus dichlorprop at 2.24 kg in 225 l: 15 Apr. Combine harvested: 24 Aug. Variety: Joss Cambier.

Permanent Grasses:

The 23rd year reseeded grass, blocks 1 and 3. PK applied: 20 Oct, 1970. NK applied to 'all-grass' half plots, and K to 'clover-grass' half plots: 17 Mar, 1971. Cut four times: 19 May, 8 July, 13 Sept, 2 Nov. NK applied to 'all-grass' half plots and K to 'clover-grass' half plots after each cut except the last.

71/R/RN/1 and 71/R/RN/2

SUMMARY OF RESULTS

WHEAT 2ND TEST CROP

GRAIN: TONNES/HECTARE

HIGHFIELD

	1967 - 69				Mean	1949-63
	LJ	LC	LN	AH		R*
Mean	6.65	6.17	6.86	6.62	6.58	6.06
1971						
NO	6.18	5.72	5.46	5.43	5.70	5.74
N1	7.24	6.32	7.35	6.71	6.90	6.07
N2	7.32	6.64	7.61	7.10	7.17	6.42
N3	5.87	6.01	7.03	7.24	6.54	5.99
1970						
F	6.74	6.22	6.78	6.70	6.61	6.51
D	6.57	6.12	6.95	6.54	6.54	5.60
1970						
NO	7.33	6.30	6.72	6.85	6.80	5.97
N1	7.03	6.61	7.32	6.89	6.96	6.50
N2	6.31	6.09	6.46	6.36	6.31	5.74
N3	5.94	5.69	6.94	6.38	6.23	6.02

Mean D.M. %: 79.8

* AH since 1964

71/R/RN/1 and 71/R/RN/2

WHEAT 2ND TEST CROP

STRAW: TONNES/HECTARE

HIGHFIELD

	1967 - 69				Mean	1949-63
	LU	LC	LN	AH		R*
Mean	7.66	7.68	7.17	6.97	7.37	7.14
1971						
NO	7.02	7.02	6.36	5.57	6.49	6.14
N1	8.56	7.94	7.44	7.11	7.76	7.28
N2	7.90	7.92	7.87	7.45	7.78	7.84
N3	7.17	7.85	7.01	7.73	7.44	7.32
1970						
F	7.54	7.59	7.09	6.82	7.26	6.99
D	7.79	7.77	7.25	7.11	7.48	7.30
1970						
NO	7.53	7.26	6.89	6.72	7.10	7.03
N1	7.84	7.74	7.39	7.09	7.51	7.03
N2	7.72	7.75	6.98	6.81	7.32	7.30
N3	7.56	7.97	7.43	7.24	7.55	7.21

Mean D.M. %: 87.6

* AH since 1964

71/R/RN/1 and 71/R/RN/2

WHEAT 2ND TEST CROP

GRAIN: TONNES/HECTARE

FOSTERS

	1967 - 69				Mean	1949-63
	LU	LC	LN	AH		R*
Mean	6.74	6.67	6.46	6.17	6.51	6.56
1971						
NO	6.03	6.09	5.83	5.04	5.75	5.31
N1	7.21	7.09	6.87	6.24	6.85	6.92
N2	7.27	7.19	7.05	6.94	7.11	7.25
N3	6.47	6.30	6.10	6.48	6.34	6.75
1970						
F	6.64	6.54	6.34	5.69	6.30	6.68
D	6.85	6.80	6.59	6.66	6.72	6.44
1970						
NO	7.00	6.62	6.47	5.91	6.50	6.48
N1	6.79	6.83	6.14	5.91	6.42	6.32
N2	6.52	6.58	6.81	6.46	6.59	6.47
N3	6.66	6.64	6.43	6.41	6.54	6.96

Mean D.M. %: 79.0

* AH since 1964

71/R/RN/1 and 71/R/RN/2

WHEAT 2ND TEST CROP

STRAW: TONNES/HECTARE

POSTERS

	1967 - 69				Mean	1949-63
	LU	LC	LN	AH		R*
Mean	7.08	7.19	6.61	6.54	6.86	7.02
1971						
NO	5.77	6.02	5.53	4.90	5.55	5.73
N1	7.07	7.49	6.69	6.28	6.88	7.07
N2	8.18	7.54	7.71	7.25	7.67	7.78
N3	7.29	7.72	6.53	7.73	7.32	7.51
1970						
F	6.71	6.76	6.17	6.03	6.42	6.77
D	7.45	7.63	7.06	7.05	7.30	7.27
1970						
NO	7.12	7.15	6.33	5.73	6.58	6.87
N1	7.07	6.98	6.39	6.21	6.66	6.71
N2	6.74	7.27	6.85	6.78	6.91	6.93
N3	7.38	7.38	6.88	7.43	7.27	7.58

Mean D.M. %: 87.3

* AH since 1964

7L/R/RN/1 and 7L/R/RN/2
 BARLEY 3RD TEST CROP
 GRAIN: TONNES/HECTARE

HIGHFIELD

	1966 - 68					1951-62		1951-68		Mean
	LU	LC	LN	AH	Mean	R*	GC	GN		
Mean	5.78	6.14	6.04	5.89	5.96	6.27	6.09	5.94	6.01	
1971										
N0	5.29	5.82	5.24	4.91	5.31	6.07	6.27	6.27	6.27	
N1	6.27	6.29	6.76	6.37	6.42	6.61	6.55	5.96	6.20	
N2	5.92	6.05	5.93	6.15	6.01	6.22	5.52	5.88	5.70	
N3	5.64	6.40	6.25	6.14	6.11	6.19	6.01	5.64	5.82	
1970										
N0	6.35	6.09	5.70	5.85	6.00					
N1	5.47	6.18	6.04	5.82	5.88					
N2	5.75	5.84	6.22	6.01	5.96					
N3	5.56	6.45	6.22	5.89	6.03					
1969										
F	5.71	6.12	6.10	5.59	5.88				6.14	
D	5.85	6.16	5.99	6.19	6.05				6.40	

Mean D.M. %: 76.3

* AH since 1963

71/R/RN/1 and 71/R/RN/2

BARLEY 3RD TEST CROP

GRAIN: TONNES/HECTARE

FOSTERS

	1966 - 68				Mean	1951-62
	LU	LC	LN	AH		R*
Mean	6.30	6.22	6.19	5.92	6.16	6.31
1971						
NO	5.19	5.85	5.41	4.64	5.27	5.32
N1	6.25	6.17	6.44	6.12	6.24	6.49
N2	7.26	6.75	6.38	6.29	6.67	6.69
N3	6.51	6.12	6.55	6.65	6.46	6.76
1970						
NO	6.21	6.21	6.21	5.31	5.98	6.65
N1	6.18	6.24	5.87	6.27	6.14	6.11
N2	6.64	6.14	6.22	5.94	6.24	6.34
N3	6.18	6.30	6.47	6.17	6.28	6.16
1969						
F	6.33	6.03	6.15	5.93	6.11	5.94
D	6.27	6.42	6.24	5.91	6.21	6.69

Mean D.M. %: 78.8

* AH since 1963

7L/R/RN/1 and 7L/R/RN/2

WHEAT 4TH TEST CROP

GRAIN: TONNES/HECTARE

HIGHFIELD

1971	1965 - 67				1950 - 67				Mean	
	LU	LC	LN	AH	Mean	RC	RN	GC		GN
N1	4.87	6.47	6.69	6.10	6.03	7.01	7.42	6.72	6.73	6.97
N2	5.17	6.76	6.57	6.32	6.21	6.35	7.08	5.83	7.35	6.65
N3	5.23	6.57	6.82	5.72	6.08	6.56	5.67	5.93	5.99	6.04
N4	5.71	5.76	5.75	5.96	5.80	5.32	5.58	4.68	5.79	5.34
Mean	5.24	6.39	6.46	6.03	6.03	6.31	6.44	5.79	6.46	6.25

Mean D.M. %: 82.1

71/R/RN/1 and 71/R/RN/2

WHEAT 4TH TEST CROP

GRAIN: TONNES/HECTARE

FOSTERS

	1965 - 67				1950 - 67			
	IU	IC	IN	AH	Mean	RC	RN	Mean
1971								
N1	6.18	6.22	6.34	5.72	6.12	7.14	7.01	7.07
N2	7.02	6.41	6.87	6.46	6.69	6.62	6.27	6.45
N3	6.57	6.51	6.42	6.49	6.50	5.82	5.72	5.77
N4	6.00	6.00	5.94	6.37	6.08	6.55	5.90	6.23
Mean	6.45	6.28	6.39	6.26	6.35	6.53	6.23	6.38

Mean D.M. %: 79.5

TL/R/RN/1 and TL/R/RN/2

WHEAT 6TH TEST CROP

GRAIN: TONNES/HECTARE

HIGHFIELD

1971	1963 - 65				Mean	1951 - 68				Mean
	IJ	IC	IN	AH		RC	RN	GC	GN	
N1	4.90	5.30	3.41	4.76	4.59	5.54	5.39	5.45	4.20	5.14
N2	5.98	5.70	4.21	5.56	5.36	4.26	5.72	6.39	4.10	5.12
N3	5.25	5.16	4.40	5.80	5.16	4.28	5.35	3.90	4.42	4.49
N4	5.70	4.53	3.94	5.70	4.97	4.26	4.39	5.28	3.94	4.47
Mean	5.46	5.17	3.99	5.45	5.02	4.58	5.21	5.26	4.16	4.80

Mean D.M. %: 84.7

71/R/RN/1 and 71/R/RN/2

WHEAT 6TH TEST CROP

GRAIN: TONNES/HECTARE

FOSTERS

1971	1963 - 65				1951 - 68	
	LU	LC	LN	AH	Mean	Mean
N1	5.42	5.94	5.40	5.61	5.59	6.11
N2	6.39	6.78	6.48	6.90	6.63	6.79
N3	6.52	6.64	6.43	6.52	6.53	6.43
N4	6.12	5.77	6.16	6.36	6.10	5.77
Mean	6.11	6.28	6.12	6.35	6.21	6.27

Mean D.M. %: 79.9

7L/R/RN/1 and 7L/R/RN/2

WHEAT 7TH TEST CROP

GRAIN: TONNES/HECTARE

HIGHFIELD

	1962 - 64				1950-64		1950 - 67		Mean
	IU	IC	IN	AH	R	Mean	GC	GN	
N1	5.09	5.17	5.54	5.83	6.51	5.63	4.07	2.86	3.46
N2	5.56	5.45	6.08	6.62	6.55	6.05	3.81	4.19	4.00
N3	5.57	6.18	6.00	6.71	6.06	6.11	4.94	3.60	4.27
N4	5.62	4.92	5.61	6.01	5.23	5.48	4.87	4.64	4.75
Mean	5.46	5.43	5.81	6.29	6.09	5.82	4.42	3.82	4.12

Mean D.M. %: 81.8

71/R/RN/1 and 71/R/RN/2

WHEAT 7TH TEST CROP

GRAIN: TONNES/HECTARE

FOSTERS

	1962 - 64				1950-64	Mean
	LU	LC	LN	AH	R	
1971						
N1	5.17	4.85	5.35	4.68	5.46	5.10
N2	6.59	6.35	6.59	6.07	5.90	6.30
N3	6.38	6.15	6.05	6.45	6.26	6.26
N4	5.78	5.73	5.68	5.82	5.77	5.76
Mean	5.98	5.77	5.92	5.75	5.85	5.85

Mean D.M. %: 78.7

71/R/RN/1 and 71/R/RN/2

HAY

DRY MATTER: TONNES/HECTARE

1st cut	2nd cut	3rd cut	4th cut	Total
HIGHFIELD				
6.50	3.01	4.13	1.86	15.50

Mean D.M. %: 1st cut: 15.9
 2nd cut: 21.6
 3rd cut: 25.2
 4th cut: 22.5
 Total of 4 cuts: 21.3

FOSTERS				
6.25	3.72	2.25	1.32	13.54

Mean D.M. %: 1st cut: 16.3
 2nd cut: 22.1
 3rd cut: 24.9
 4th cut: 22.3
 Total of 4 cuts: 21.4

71/R/RN/1 and 71/R/RN/2

	HIGHFIELD	FOSTERS
	Mean	Mean

LUCERNE, DRY MATTER: TONNES/HECTARE

TOTAL OF 3 CUTS

2nd year	4.66	5.56
----------	------	------

ALL-GRASS LEY, DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

2nd year	7.33	6.79
----------	------	------

CLOVER-GRASS LEY, DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

2nd year	3.22	4.06
----------	------	------

RESEEDED GRASS, DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

	HIGHFIELD			FOSTERS		
	Blocks	RC	RN	Blocks	RC	RN
23rd Exptl year	1 & 4	3.46	11.17	1 & 4	5.90	9.56

71/R/RN/1 and 71/R/RN/2

PERMANENT GRASS, DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

	GC	GN
HIGHFIELD		
23rd Exptl year		
Blocks 1 & 4	3.21	10.41
Block 2	2.97	10.24

(C) Clover-grass management
(N) All-grass management