

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1971

[Full Table of Content](#)



### 71/R/RN/5 Arable Reference Plots - Barley, Ley, Potatoes, Wheat, Kale

#### Rothamsted Research

Rothamsted Research (1972) *71/R/RN/5 Arable Reference Plots - Barley, Ley, Potatoes, Wheat, Kale ; Yields Of The Field Experiments 1971*, pp 80 - 83 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/RN/5

ARABLE REFERENCE PLOTS

Object: To study the long term effects of FYM and N, P and K fertilisers on the yield and mineral content of crops - Great Field IV.

The sixteenth year of the rotation barley, ley, potatoes, winter wheat, kale. The twelfth year of the same rotation on the additional plots. The fifteenth year of permanent grass.

For previous years see 58/Bc/1(t), 59/Bc/1(t), 60/B/3(t), 61-64/B/2, 65/B/2(t), 66/B/2(t), 67/B/2, 68/B/3(t) and 69-70/R/RN/5.

Whole plot dimensions: 2.13 x 2.44.

Cultivations, etc.: Ground chalk applied at 3.76 tonnes to all plots of the main experiment: 27 Nov, 1970.

Winter wheat: Balancing Mg applied to half plots: 25 Sept, 1970.

Plots dug by hand, test Mg applied: 2 Oct. P, K, Ca and S applied, seed drilled: 5 Oct. First half N dressing applied: 9 Mar, 1971. Trace element spray applied: 19 Apr. All N applied to additional plots, second half N dressing applied: 27 Apr. Harvested: 13 Aug. Variety: Maris Nimrod.

Kale: FYM applied, plots dug by hand: 3 Nov, 1970. P, K, Ca, Mg and S applied to additional plots: 29 Jan, 1971. P and K applied to remainder: 19 Feb. First half N dressing applied to additional plots, all N to remainder, plots rotary cultivated, seed drilled: 7 Apr. Trace element spray applied: 11 June. Second half N dressing applied to additional plots: 15 June. Sprayed with menazon ('Saphicol' at 0.7 l in 450 l): 6 July. Harvested: 22 Oct. Variety: Thousand headed.

Barley: Plots dug by hand: 3 Nov, 1970. P, K, Mg, Ca and S applied to additional plots: 29 Jan, 1971. P and K applied to remainder: 19 Feb. N applied, plots rotary cultivated, seed drilled: 31 Mar. Trace element spray applied: 24 May. Sprayed (excluding additional plots) with tridemorph fungicide at 0.53 kg in 450 l): 15 June. Harvested - additional plots: 13 Aug, remainder: 16 Aug. Variety: Deba Abed (Midas on additional plots, seed dressed with ethirimol).

Grass-clover ley: Seed drilled in barley stubble: 14 Aug, 1970. P, K, Ca, Mg and S applied: 17 Dec. N applied: 9 Mar, 1971. Trace element spray applied: 19 Apr. Cut three times: 4 June, 23 July, 20 Sept. Varieties: R.V.P. Italian Ryegrass and Dorset Marl Clover.

71/R/RN/5

Potatoes: FYM applied and plots dug by hand: 4 Nov, 1970. P, K, Ca, Mg and S applied to additional plots: 29 Jan, 1971. P and K applied to remainder: 19 Feb. First half N dressing applied to additional plots, all N to remainder, plots rotary cultivated, Mg applied to half plots, potatoes planted: 7 Apr. Trace element spray applied: 11 June. Second half N dressing applied to additional plots: 15 June. All plots sprayed on 2 occasions with mancozeb at 1.35 kg plus menazon ('Saphicol' at 0.7 l) in 450 l: 16 June, 28 July. Additional plots only sprayed once as above described: 6 July. Lifted: Plots of main experiment with neither K nor FYM and no fertiliser plots of additional plots: 30 July. Remainder sprayed with captafol fungicide at 1.68 kg plus menazon ('Saphicol' at 0.7 l) in 450 l: 18 Aug. Remaining plots lifted: 14 Sept. Variety: King Edward.

Permanent grass: P and K applied: 17 Dec, 1970. FYM applied: 22 Feb, 1971. N applied: 9 Mar, 20 May, 15 July. Cut three times: 20 May, 15 July, 11 Oct.

- NOTES: (1) Yields of dry matter were obtained for each crop.  
(2) The percentages of N, P and K were measured in each crop.  
(3) The percentage of Mg was measured in potato tubers on the main experiment.  
(4) The percentage of K in potato leaves was measured on the main experiment.



71/R/RN/5

SUMMARY OF RESULTS

GREAT FIELD IV (R): ORIGINAL PLOTS

TONNES/HECTARE

Treatment	WINTER WHEAT:		KALE:		BARLEY:		LEY: DRY MATTER			POTATOES		PERMANENT GRASS:		
	GRAIN	STRAW	FRESH WEIGHT	GRAIN	STRAW	1st cut	2nd cut	3rd cut	Total of 3 cuts	TOTAL TUBERS	1st cut	2nd cut	3rd cut	Total of 3 cuts
O	3.22	5.20	15.7	2.25	2.36	2.19	0.81	0.97	3.97	7.1	0.73	1.17	1.00	2.90
N1	1.34	4.62	20.9	2.73	3.42	5.16	1.20	1.34	7.70	8.1	1.14	1.39	1.41	3.94
P	2.78	4.34	27.5	3.40	2.87	2.86	0.67	0.50	4.03	13.4	0.59	0.94	0.67	2.20
N1P	0.88	3.56	37.5	2.56	3.28	5.23	0.93	0.36	6.52	8.0	1.75	1.50	1.66	4.91
K	4.18	5.68	12.6	3.11	2.58	3.72	1.95	1.83	7.50	33.8	0.76	1.05	1.11	2.92
N1K	5.69	7.53	13.1	1.53	2.98	4.78	1.86	2.29	8.93	37.1	2.12	1.90	2.03	6.05
PK	4.41	5.93	27.5	3.76	2.79	4.74	3.43	3.26	11.43	36.3	1.00	1.74	1.17	3.91
N1PK	6.52	9.34	42.7	4.79	5.24	6.35	2.71	2.81	11.87	43.4	2.39	1.98	1.60	5.97
N2PK	8.50	11.48	48.4	5.15	5.55	7.40	2.18	2.20	11.78	51.0	4.12	2.49	2.55	9.16
D	5.02	7.74	34.4	4.49	3.81	5.97	2.29	2.26	10.52	45.9	3.75	1.67	1.54	6.96
N1PKD	7.44	11.75	53.2	5.84	5.80	6.45	2.26	2.26	10.97	58.1	3.68	2.59	2.38	8.65
N2PKD	5.00	13.33	61.0	5.55	7.31	7.72	2.09	1.93	11.74	64.9	4.50	4.33	2.43	11.26
Mean	80.0	65.1		74.7	61.2	27.3	25.8	29.6	27.6		27.8	29.2	33.0	30.0
D.M.%:														

71/R/RW/5

GREAT FIELD IV (R): ADDITIONAL PLOTS

TONNES/HECTARE

Treatment	WINTER WHEAT:		KALE: FRESH WEIGHT	BARLEY:		LEY: DRY MATTER			Total of 3 cuts	POTATOES TOTAL TUBERS
	GRAIN	STRAW		GRAIN	STRAW	1st cut	2nd cut	3rd cut		
None	3.33	4.73	17.9	2.86	3.36	2.84	0.76	1.02	4.62	7.0
N2 PK	6.38	7.52	57.6	5.87	7.49	7.31	2.28	2.12	11.71	41.8
N2 PK Mg Ca	7.42	8.67	60.2	4.94	7.93	7.39	2.56	1.86	11.81	32.3
N2 PK Mg S	7.38	8.74	59.7	4.89	6.94	6.65	1.76	1.73	10.14	38.4
N2 PK Ca S	7.39	9.25	59.7	5.29	7.39	7.45	2.13	1.98	11.56	34.4
N2 PK Mg Ca S	7.48	9.22	60.6	6.07	7.46	8.91	2.85	2.46	14.22	34.0
N2 PK Mg Ca S TE	6.66	8.84	61.5	5.81	7.34	7.74	2.31	2.24	12.29	31.4
Mean D.M. %:	80.2	67.5		81.1	78.3	28.5	26.1	28.8	27.8	