


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1971

[Full Table of Content](#)



## 71/R/RN/7 Residual Phosphate - Potatoes, Barley, Swedes

### Rothamsted Research

Rothamsted Research (1972) *71/R/RN/7 Residual Phosphate - Potatoes, Barley, Swedes ; Yields Of The Field Experiments 1971*, pp 88 - 91 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/RN/7

RESIDUAL PHOSPHATE

Object: To study direct and residual effects of phosphate fertiliser on yields of three crops grown in rotation - Sawyers I and Great Field IV.

The 12th year, potatoes, barley, swedes.

For previous years see 'Details' 1967 and 68/B/5(t), 69/R/RN/7 and 70/R/RN/7(t).

Whole plot dimensions:-

Sawyers I: 4.27 x 20.1. Area harvested: Potatoes and barley - 0.00572, Swedes - 0.00429.

Great Field IV: 4.27 x 19.3. Area harvested: Potatoes and barley - 0.00520, Swedes - 0.00390.

Standard applications:

Potatoes: Weedkillers: Linuron at 0.84 kg plus paraquat at 0.42 kg in 427 l. Fungicide: Mancozeb at 1.34 kg in 438 l on 2 occasions. Insecticide: Demeton-s-methyl at 245 g applied with the fungicide on the first occasion.

Barley: Ground chalk at 3140 kg. Weedkiller: Ioxynil octanoate, bromoxynil octanoate and the iso-octyl ester of dichlorprop ('Oxytril P' at 1.4 l in 225 l).

Cultivations, etc. (both fields):- Ploughed: 27 Oct, 1970.

Potatoes: Fertilisers applied: 6 Apr, 1971. Plots rotary cultivated, potatoes machine planted: 7 Apr. Weedkiller applied: 10 May. Grubbed: 5 June. Rotary ridged: 7 June. Fungicide plus insecticide applied: 24 June. Fungicide applied: 13 Aug. Sprayed with undiluted BOV at 168 l: 15 Sept. Lifted: 24 Sept. Variety: Majestic.

Barley: Chalk applied: 12 Oct, 1970. Treatment P fertilisers applied: 3 Mar, 1971. Seed combine drilled at 157 kg: 10 Mar. Weedkiller applied: 4 May. Combine harvested: 17 Aug. Variety: Julia.

Swedes: Treatment P and basal K applied: 21 Apr, 1970. N applied: 14 May. Seed drilled at 1.7 kg: 18 May. Singled: 28 June. Lifted: 20 Oct. Variety: Wilhelmsburger.

71/R/RN/7

Standard errors per plot.

Sawyers I:

Potatoes, total tubers, tonnes/hectare:	2.11 or 6.7% (11 d.f.)
Barley, grain, tonnes/hectare:	0.224 or 4.0% (11 d.f.)
Swedes, roots, tonnes/hectare:	2.30 or 10.4% (11 d.f.)

71/R/RN/7

SUMMARY OF RESULTS

POTATOES

Treatment	TOTAL TUBERS: TONNES/HECTARE % WARE: 3.81 CM (1.5 INCH)			
	Great Field IV	Sawyers I	Great Field IV	Sawyers I
		(±1.49)		
O	27.9	23.6	96.6	97.3
A1	32.1	31.8	94.8	95.0
A2	34.3	36.9	94.3	93.1
A3	40.2	42.6	94.6	93.2
A4	46.3	42.3	93.4	93.6
T1	28.1	26.3	96.1	95.5
T2	31.7	28.0	93.8	93.6
R2	34.5	29.4	95.6	95.2
R3	29.5	31.0	93.0	93.6
R4	39.8	38.8	93.7	93.1
G1	28.9	23.8	94.8	95.5
S1	27.8	22.6	95.7	96.2
Mean	33.4	31.4	94.7	94.6

BARLEY

	GRAIN: TONNES/HECTARE		STRAW: TONNES/HECTARE	
			(±0.158)	
O	3.58	4.90	2.98	3.12
A1	4.72	5.70	4.54	3.71
A2	4.60	5.78	3.82	3.79
A3	4.56	5.95	4.01	4.08
A4	5.06	6.28	4.17	4.21
T1	3.94	5.62	3.98	3.69
T2	4.02	5.64	2.86	3.79
R2	4.47	5.41	3.48	3.26
R3	4.60	5.77	4.02	3.94
R4	4.40	5.87	3.67	4.18
G1	4.55	5.35	3.59	3.76
S1	3.09	5.19	3.54	3.56
Mean	4.30	5.62	3.72	3.76
Mean D.M. %:	82.4	83.8	83.8	87.2

71/R/RN/7

SWEDES, ROOTS: TONNES/HECTARE

Treat- ment	Great Field IV	Sawyers I
		(±1.63)
0	10.7	9.3
A1	22.0	23.5
A2	28.5	28.0
A3	30.6	27.4
A4	25.5	29.3
T1	19.1	19.0
T2	23.9	21.5
R2	23.8	21.5
R3	26.4	28.9
R4	30.7	28.7
G1	16.0	13.7
S1	13.4	15.5
Mean	22.5	22.2