

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/W/RN/13 Intensive Cereals - Ley, Potatoes, W. Wheat, Barley

**Rothamsted Research**

Rothamsted Research (1973) 72/W/RN/13 *Intensive Cereals - Ley, Potatoes, W. Wheat, Barley* ; Yields Of The Field Experiments 1972, pp 114 - 118 - DOI:

<https://doi.org/10.23637/ERADOC-1-62>

72/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne diseases and organic matter in the soil - Woburn Stackyard I.

Sponsors: G.W. Cooke, D.B. Slope.

The seventh year, ley, potatoes, winter wheat, barley.

For previous years see 66/B/9(t), 67/B/9, 68/B/7(t), 69/W/RN/13(t), 70/W/RN/13(t) and 71/W/RN/13(t).

Whole plot dimensions: 8.53 x 20.4. Sub plot area harvested:  
Potatoes - 0.00139, wheat - 0.00273, barley - 0.00273.

Basal applications: All crops: 130 kg P2O5, 260 kg K2O as (0:14:28), half ploughed in, half applied to the plough furrow. Weedkiller to all crops except potatoes: Paraquat at 0.56 kg ion in 280 l.  
Leys: 63 kg N in seedbed, 63 kg N after sowing as 'Nitro-Chalk'. Only one cut was taken.

Potatoes: 150 kg N as 'Nitro-Chalk'. Weedkiller: Linuron at 1.1 kg with paraquat at 0.4 kg ion in 370 l. Fungicide: Mancozeb at 1.3 kg in 430 l. Fungicide with insecticide: Mancozeb at 1.3 kg with demeton-s-methyl at 0.25 kg in 430 l.

Wheat: Weedkiller: 2,4-D at 0.7 kg with dichlorprop at 2.8 kg in 280 l.

Barley: Weedkiller: 2,4-D at 0.56 kg with dichlorprop at 2.2 kg in 280 l.

Seed: Leys: Italian ryegrass S22, sown at 28 kg.

Potatoes: Majestic.

Wheat: Cappelle, sown at 180 kg.

Barley: Julia (dressed with ethirimol) sown at 180 kg.

Cultivations, etc.: All plots: Half PK applied, ploughed: 1 Oct, 1971. Remaining PK applied: 7 Oct.

Leys: Paraquat applied: 9 Sept, 1971. Deep-tine cultivated, wheat block only: 4 Nov. N applied: 18 Apr, 1972. Rotary cultivated: 21 Apr. Seed sown: 26 Apr. N applied: 6 June. Cut once: 6 Sept.

72/W/RN/13

Potatoes: Deep-tine cultivated, wheat block only: 4 Nov, 1971.  
N applied: 4 Apr, 1972. Rotary cultivated, potatoes planted:  
18 Apr. Linuron with paraquat applied: 6 May. Rotary ridged:  
19 June. Fungicide applied: 5 July. Fungicide with insecticide  
applied: 27 July. Haulm mechanically destroyed: 14 Sept.  
Lifted: 20 Sept.

Wheat: Paraquat applied: 9 Sept, 1971. Seed drilled: 15 Oct. N  
applied: 11 Apr, 1972. 2,4-D with dichlorprop applied: 27 Apr.  
Combine harvested: 30 Aug.

Barley: Paraquat applied: 9 Sept, 1971. Seed drilled: 15 Mar, 1972.  
N applied: 16 Mar. 2,4-D with dichlorprop applied: 8 May.  
Combine harvested: 24 Aug.

NOTE: Estimates of eyespot (*Cercosporaella herpotrichoides*) and take-all  
(*Gaeumannomyces graminis* formerly *Ophiobolus graminis*) were  
made on both cereal crops.

Standard errors per sub plot.

Wheat, grain, tonnes/hectare: 0.389 or 16.7% (12 d.f.)  
Barley, grain, tonnes/hectare: 0.197 or 4.0% (11 d.f.)

NOTES: (1) One barley plot (CS6, NSO) was severely damaged by birds. The  
yield of this plot was not recorded - an estimated value  
was used in the analysis.  
(2) In 1972 and in 1971 the yields of leys were not recorded.

72/W/RN/13

TABLES OF MEANS

POTATOES

N: KG/Ha 1970

63	126	189	252	Mean
----	-----	-----	-----	------

PERMANENT WHEAT BLOCKS

TOTAL TUBERS: TONNES/HECTARE

32.9	38.1	35.6	36.2	35.7
------	------	------	------	------

PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE

93.1	94.0	94.5	93.4	93.7
------	------	------	------	------

PERMANENT BARLEY BLOCKS

TOTAL TUBERS: TONNES/HECTARE

41.7	40.8	39.4	43.9	41.4
------	------	------	------	------

PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE

95.0	94.8	94.8	95.3	95.0
------	------	------	------	------

72/W/RN/13

WINTER WHEAT

N: KG/HA

CROP SEQUENCE (CS)		63	126	189	252	Mean
GRAIN: TONNES/HECTARE						
1966	1967	1968	1969	1970	1971	
2 P	W	W	W	L	P	2.10
3 W	W	W	L	P	W	1.37
4 W	W	L	P	W	W	1.02
6 W	W	W	W	W	W	1.16
						3.55
						3.39
						3.81
						3.21
						2.27
						1.95
						1.80
Mean			1.41	2.49	2.74	2.69
2.33						

STANDARD ERRORS OF DIFFERENCES  
N CS\* N

0.194 0.389

\* Within the same level of CS only

STRAW: TONNES/HECTARE						
1966 1967 1968 1969 1970 1971						
2	P	W	W	W	L	P
2 P	W	W	W	L	P	W
3 W	W	W	L	P	W	1.65
4 W	W	L	P	W	W	1.78
6 W	W	W	W	W	W	3.02
						4.21
						3.36
						3.34
						3.54
						3.02
						2.85
						3.61
						3.56
						3.61
Mean			2.13	3.49	3.58	3.76
3.24						

Mean D.M. %: Grain: 82.6  
Straw: 87.0

72/W/RN/13

BARLEY

N: KG/HA

CROP SEQUENCE (CS)						50	100	150	200	Mean
GRAIN: TONNES/HECTARE										
2	P	B	B	B	L	P	3.83	5.42	6.01	5.78
3	B	B	B	L	P	B	3.68	5.33	6.01	5.78
4	B	B	L	P	B	B	3.42	4.99	5.71	5.98
6	B	B	B	B	B	B	1.92	4.46	5.40	5.22
Mean						3.21	5.05	5.78	5.69	4.93

STANDARD ERRORS OF DIFFERENCES

N	CS*	N
0.098	0.197	

\* Within the same level of CS only

STRAW: TONNES/HECTARE

1966 1967 1968 1969 1970 1971						3.05	4.63	5.41	5.28	4.59
2	P	B	B	B	L	P	2.80	3.82	4.64	5.10
3	B	B	B	L	P	B	2.78	3.77	4.71	5.21
4	B	B	L	P	B	B	2.43	3.70	5.00	5.15
6	B	B	B	B	B	B				4.07
Mean						2.76	3.98	4.94	5.19	4.22

Mean D.M. %: Grain: 85.6  
Straw: 84.9