

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/CS/13 N Levels to Old Grass - Old Grass

Rothamsted Research

Rothamsted Research (1973) *72/R/CS/13 N Levels to Old Grass - Old Grass* ; Yields Of The Field Experiments 1972, pp 143 - 146 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/CS/13

N LEVELS TO OLD GRASS

Object: To study the effects of a range of nitrogen rates on yield and botanical composition of very old permanent pasture given a single dressing of P and K annually. Nitrogen fixed by legumes is estimated and the effects of treatments on nutrients available in the soil is also studied - Park Grass old plot 6.

Sponsors: A.E. Johnston, R.C. Flint.

The eighth year, old grass.

For previous years see 65/C/33(t), 66/C/14, 67/C/10(t), 68/C/8(t), 69/R/CS/13(t), 70/R/CS/13(t) and 71/R/CS/13.

Design: 4 randomised blocks of 10 plots.

Whole plot dimensions: 1.83 x 10.1. **Area harvested:** 0.00086.

Treatments:

Herbicide: Sprayed with mecoprop ('Clovotox' at 11.2 l in 450 l) to control legumes (two plots per block) (OS) - nitrogen fertiliser is not applied to this treatment.

Nitrogen (total applied per year in 4 equal dressings): None (two plots per block) (0), 75 (1), 150 (2), 225 (3), 300 (4), 375 (5), 450 (6) kg N as 'Nitro-Chalk'.

Rates 1 and 2 were previously 1 (145 kg N).

Rates 3 and 4 were previously 2 (290 kg N).

Rates 5 and 6 were previously 3 (435 kg N).

Basal applications: 34 kg P as superphosphate, 224 kg K as potassium sulphate, 11 kg Mg as magnesium sulphate.

Cultivations, etc.: Basal P K Mg applied: 6 Dec, 1971. N applied: 24 Mar, 1972. Herbicide applied to 3 plots: 27 Apr, 24 July. Plots cut: 15 May, 29 June, 14 Aug, 23 Oct. N applied after each cut except the last.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut: 0.300 or 10.3% (29 d.f.)

2nd cut: 0.232 or 11.9% (29 d.f.)

3rd cut: 0.137 or 10.6% (29 d.f.)

4th cut: 0.120 or 21.5% (29 d.f.)

Total of 4 cuts: 0.465 or 6.9% (29 d.f.)

72/R/CS/13

TABLES OF MEANS

DRY MATTER: TONNES/HECTARE

								N	
OS*	O*	1	2	3	4	5	6	Mean	
								1ST CUT	
0.60	1.66	2.17	2.80	4.38	4.77	5.00	5.40	2.90	

STANDARD ERRORS OF DIFFERENCES

Between any of N1-N6	0.212
Between NOS and NO	0.150
Between NOS or NO and any of N1-N6	0.184

								2ND CUT	
0.74	1.77	1.85	2.27	2.37	2.98	2.41	2.59	1.95	

STANDARD ERRORS OF DIFFERENCES

Between any of N1-N6	0.164
Between NOS and NO	0.116
Between NOS or NO and any of N1-N6	0.142

* Duplicated treatment

Mean D.M.%: 1st cut: 21.4
2nd cut: 24.4

72/R/CS/13

DRY MATTER: TONNES/HECTARE

N

OS*	0*	1	2	3	4	5	6	Mean
3RD CUT								
0.23	1.16	0.94	1.27	1.78	1.91	2.11	2.21	1.30

STANDARD ERRORS OF DIFFERENCES

Between any of N1-N6	0.097
Between NOS and NO	0.069
Between NOS or NO and any of N1-N6	0.084

4TH CUT

0.15	0.23	0.38	0.71	0.84	1.01	0.91	0.99	0.56
------	------	------	------	------	------	------	------	------

STANDARD ERRORS OF DIFFERENCES

Between any of N1-N6	0.085
Between NOS and NO	0.060
Between NOS or NO and any of N1-N6	0.074

* Duplicated treatment

Mean D.M.%: 3rd cut: 25.4
4th cut: 34.5

72/R/CS/13

DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

N								
OS*	0*	1	2	3	4	5	6	Mean
1.72	4.82	5.34	7.05	9.36	10.68	10.42	11.19	6.71

STANDARD ERRORS OF DIFFERENCES

Between any of N1-N6	0.329
Between NOS and NO	0.233
Between NOS or NO and any of N1-N6	0.285

* Duplicated treatment

Mean D.M.%: Total of 4 cuts: 26.4