

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/W/CS/33 Rates of Nematicides Dosage - Potatoes

Rothamsted Research

Rothamsted Research (1973) *72/W/CS/33 Rates of Nematicides Dosage - Potatoes* ; Yields Of The Field Experiments 1972, pp 176 - 178 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/W/CS/33

RATES OF NEMATOCIDES DOSAGE

Object: To study the residual effects on yield of potatoes of several nematocides applied at a range of rates (to potatoes 1969, sugar beet 1970, none to barley 1971) - Woburn Butt Close.

Sponsor: A.G. Whitehead.

The fourth year, potatoes.

For previous years see 69/W/CS/33(t), 70/W/CS/33(t) and 71/W/CS/33.

Whole plot dimensions: 2.84 x 9.14. Area harvested: 0.00065.

Basal applications: 1500 kg (13:13:20). Weedkillers: Paraquat at 0.56 kg ion in 280 l. Linuron at 1.1 kg with paraquat at 0.42 kg ion 370 l. Fungicide: Mancozeb at 1.3 kg in 430 l. Fungicide with insecticide: Mancozeb at 1.3 kg with demeton-s-methyl at 0.24 kg in 430 l.

Variety: Pentland Crown.

Cultivations, etc.: Paraquat applied: 9 Sept, 1971. Ploughed twice: 21 Sept, 17 Feb, 1972. NPK applied: 31 Mar. Rotary cultivated, potatoes planted: 5 Apr. Weedkiller applied: 5 May. Fungicide applied: 5 July. Fungicide with insecticide applied: 26 July. Haulm mechanically destroyed: 6 Sept. Lifted: 20 - 21 Sept.

NOTE: Soil samples were taken after harvest for cyst and egg counts of *Heterodera rostochiensis*.

Standard errors per plot. Total tubers, tonnes/hectare.

Whole plot: 4.59 or 15.2% (26 d.f.)

Sub plot: 4.03 or 13.3% (29 d.f.)

72/W/CS/33

TABLES OF MEANS

TOTAL TUBERS: TONNES/HECTARE

NEMATOCIDE

APPLICATION	O	M	D1	D2	D3	D4	T1	T2	T3	T4	Z1	Z2	Z3	Z4	Mean	
1969			33.6	31.4	19.1	21.7	30.0	27.0	17.4	32.3	27.8	22.4	35.7	31.7	32.8	27.9
1969-70			42.5	33.6	27.4	34.5	32.4	33.9	32.8	28.5	37.1	23.7	33.0	33.0	36.7	33.0
Mean	27.8	38.1	32.5	23.2	28.1	31.2	30.5	25.1	30.4	32.4	23.1	34.4	32.4	34.8	30.3*	

* Grand mean

STANDARD ERRORS OF DIFFERENCES

NEMATOCIDE	APPLICATION	NEMATOCIDE	APPLICATION
		3.75	0.91
Unless same level of NEMATOCIDE			4.41
			3.29

72/W/CS/33

PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE

NEMATOCIDE

APPLICATION	O	M	D1	D2	D3	D4	T1	T2	T3	T4	Z1	Z2	Z3	Z4	Mean
1969		94.5	94.0	87.9	91.1	92.8	91.2	87.7	93.7	93.3	90.7	93.2	92.1	92.0	91.9
1969-70		94.0	93.4	88.2	93.5	91.0	91.6	92.8	93.5	94.0	92.1	91.8	93.5	93.0	92.5
Mean	91.7	94.3	93.7	88.0	92.3	91.9	91.4	90.3	93.6	93.6	91.4	92.5	92.8	92.5	92.1*

* Grand mean