

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/W/CS/56 Nematodes and Verticillium - W. Wheat

### Rothamsted Research

Rothamsted Research (1973) *72/W/CS/56 Nematodes and Verticillium - W. Wheat* ; Yields Of The Field Experiments 1972, pp 204 - 205 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/W/CS/56

NEMATODES AND VERTICILLIUM

Object: To study the residual effects of methyl bromide (applied 1969 and 1971) and aldicarb and benomyl (applied 1969 only) (C) on winter wheat - Woburn Broad Mead I.

Sponsors: D.C.M. Corbett, G.A. Hide.

The fourth year, winter wheat.

For previous years see 69/W/P/2(t), 70/W/CS/56(t) and 71/W/CS/56(t).

Design: 4 blocks of 6 plots.

Whole plot dimensions: 2.67 x 13.4. Area harvested: 0.00191.

Basal applications: 290 kg (0:20:20) combine drilled, 440 kg 'Nitro-Chalk'. Weedkiller: 2,4-D at 0.70 kg plus dichlorprop at 2.8 kg in 280 l.

Seed: Cappelle sown at 170 kg..

Cultivations, etc.: Deep-tine cultivated on 2 occasions: 16 Sept, 9 Oct, 1971. Seed combine drilled: 15 Oct. Weedkiller applied: 31 Mar, 1972. N applied: 17 Apr. Combine harvested: 1 Sept.

NOTE: Grain samples were taken for analysis of methyl bromide residues.

Standard error per plot.

Grain, tonnes/hectare: 0.435 or 6.1% (15 d.f.)

72/W/CS/56

TABLE OF MEANS

GRAIN: TONNES/HECTARE

C

00	MD	MW	TO	M1	M2	Mean
7.34	7.26	7.32	6.82	7.24	7.13	7.19

STANDARD ERROR OF DIFFERENCES

C 0.308

Mean D.M. %: 85.0