

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/W/CS/63 Nematodes and Verticillium - W. Wheat

### Rothamsted Research

Rothamsted Research (1973) *72/W/CS/63 Nematodes and Verticillium - W. Wheat* ; Yields Of The Field Experiments 1972, pp 216 - 217 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/W/CS/63

NEMATODES AND VERTICILLIUM

Object: To study the residual effects of methyl bromide, aldicarb and benomyl (applied 1970) on winter wheat - Woburn Broad Mead I.

The third year, winter wheat.

Sponsors: D.C.M. Corbett, G.A. Hide.

For previous years see 70/W/P/2(t) and 71/W/CS/63.

Design: 4 blocks of 4 plots.

Whole plot dimensions: 2.67 x 13.4. Area harvested: 0.00191.

Treatments:-

Residues of chemicals applied to potatoes 1970, (none in 1971 or 1972), (C). None (O), methyl bromide at 975 kg (M), aldicarb at 7 kg (T), benomyl at 22 kg (B).

Basal applications: 290 kg (0:20:20) combine drilled, 440 kg 'Nitro-Chalk'. Weedkiller: 2,4-D at 0.7 kg plus dichlorprop at 2.8 kg in 280 l.

Seed: Cappelle sown at 170 kg.

Cultivations, etc.:

Deep-tine cultivated on two occasions: 16 Sept, 9 Oct, 1971.

Seed combine drilled: 15 Oct. Weedkiller applied: 31 Mar, 1972.

N applied: 17 Apr. Combine harvested: 1 Sept.

NOTE: Grain samples were taken for determination of residues of methyl bromide.

Standard error per plot.

Grain, tonnes/hectare: 0.272 or 3.5% (9 d.f.)

72/W/CS/63

TABLE OF MEANS

GRAIN: TONNES/HECTARE

C

O	B	M	T	Mean
7.71	7.83	7.62	7.75	7.73

STANDARD ERROR OF DIFFERENCES

C 0.193

Mean D.M. %: 84.6