

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/CS/76 and 72/W/CS/76 Rates of NPK Fertilisers - W. Wheat

Rothamsted Research

Rothamsted Research (1973) *72/R/CS/76 and 72/W/CS/76 Rates of NPK Fertilisers - W. Wheat ; Yields Of The Field Experiments 1972*, pp 222 - 225 - DOI:

<https://doi.org/10.23637/ERADOC-1-62>

72/R/CS/76 and 72/W/CS/76

RATES OF NPK FERTILISER

Object: To study the residual effects of a range of rates of compound NPK fertiliser, applied to potatoes in 1971, on the yield of wheat. The direct effects of nitrogen are also studied - Rothamsted (R) Long Hoos III and Woburn (W) Lansome II.

Sponsor: F.V. Widdowson.

The second year, winter wheat.

For previous year see 71/R&W/CS/76(t).

Design: 4 x 2 x 2 x 2 in 2 blocks of 8 plots split into 2.

Whole plot dimensions:

Long Hoos III (R) and Lansome II (W): 4.27 x 21.0. Sub plot area harvested: 0.00286.

Treatments: All combinations of:-

Whole plots: 1. Rates of compound fertiliser (13:13:20) to potatoes in 1971: 1260, 1880, 2510, 3140 kg.

2. Spacing of potatoes within the rows in 1971: 30 cm (12 inches), 45 cm (18 inches).

Sub plots: 3. Variety of potatoes in 1971:

Long Hoos III (R): King Edward (KE), Pentland Crown (PC).

Lansome II (W): Pentland Crown (PC), Record (R).

4. Nitrogen to wheat in 1972:

Long Hoos III (R): 38, 76 kg N as 'Nitro-Chalk'.

Lansome II (W): 63, 126 kg N as 'Nitro-Chalk'.

Basal applications:

Long Hoos III (R): Manures: None. Weedkillers: Ioxynil at 0.63 kg plus mecoprop at 1.89 kg in 220 l.

Lansome II (W): 2,4-D at 0.70 kg plus dichlorprop at 2.8 kg in 280 l.

Seed: Cappelle sown at 200 kg (R), 180 kg (W).

72/R/CS/76 and 72/W/CS/76

TABLES OF MEANS

LONG HODS III (R)

COMPOUND FERTILISER (13:13:20) KG/HA (F)

	1260	1880	2510	3140	Mean
GRAIN: TONNES/HECTARE					
SPACING: CM					
30	7.07	7.08	7.39	7.43	7.24
45	6.98	7.23	7.19	7.43	7.21
VARIETY					
KE	7.10	7.21	7.29	7.24	7.21
PC	6.95	7.11	7.30	7.61	7.24
N: KG/HA					
38	6.76	6.90	6.99	7.35	7.00
76	7.30	7.41	7.60	7.50	7.45
Mean	7.03	7.16	7.29	7.43	7.23

STANDARD ERRORS OF DIFFERENCES

F	SPACING	VARIETY	N
0.173	0.122	0.087	0.087
F	F	F	
SPACING	VARIETY	N	
0.244	0.212	0.212	
Unless same level of F		0.174	0.174

Mean D.M. %: 85.5

72/R/CS/76 and 72/W/CS/76

Cultivations, etc.:

Long Hoos III (R): Deep-tine cultivated twice: 6 Oct, 1971. Seed drilled: 11 Oct. N applied: 6 Apr, 1972. Weedkiller applied: 28 Apr. Combine harvested: 31 Aug.

Lansome II (W): Deep-tine cultivated on 2 occasions: 6 Oct and 9 Oct, 1971. Seed drilled: 22 Oct. N applied: 12 Apr, 1972. Weedkiller applied: 27 Apr. Combine harvested: 31 Aug.

NOTE: The percentage of N in the grain was determined.

Standard errors per plot. Grain, tonnes/hectare:

Long Hoos III (R): Whole plot: 0.244 or 3.4% (6 d.f.)

Sub plot: 0.247 or 3.4% (6 d.f.)

Lansome II (W): Whole plot: 0.314 or 7.4% (6 d.f.)

Sub plot: 0.421 or 9.9% (6 d.f.)

72/R/CS/76 and 72/W/CS/76

LANSOME II (W)

COMPOUND FERTILISER (13:13:20) KG/HA (F)

	1260	1880	2510	3140	Mean
SPACING: CM	GRAIN: TONNES/HECTARE				
30	4.36	4.23	4.25	3.99	4.21
45	4.25	4.18	4.07	4.62	4.28
VARIETY					
PC	4.26	4.18	4.20	4.49	4.28
R	4.36	4.23	4.12	4.13	4.21
N: KG/HA					
63	3.79	3.72	3.82	3.97	3.83
126	4.82	4.69	4.50	4.65	4.67
Mean	4.31	4.21	4.16	4.31	4.25

STANDARD ERRORS OF DIFFERENCES

F	SPACING	VARIETY	N
0.222	0.157	0.149	0.149

F	F	F
SPACING	VARIETY	N

Unless same level of
F

0.314 0.306 0.306

0.298 0.298

Mean D.M. %: 81.9