

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/W/CS/77 Much Fertiliser and Fym - W. Wheat

#### Rothamsted Research

Rothamsted Research (1973) *72/W/CS/77 Much Fertiliser and Fym - W. Wheat* ; Yields Of The Field Experiments 1972, pp 226 - 228 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/W/CS/77

MUCH FERTILISER AND FYM

Object: To study the residual effects of large dressings of FYM and NPK fertiliser applied to potatoes 1971, on the yields of a following winter wheat - Woburn Stackyard C.

Sponsor: F.V. Widdowson.

The second year, winter wheat.

For previous year see 71/W/CS/77(t).

Design: 3 blocks of 12 plots split into 2.

Whole plot dimensions: 2.74 x 3.05. Area harvested: 0.00033.

Treatments: All combinations of:-

Whole plots: 1. Residues of FYM, fertiliser and methods of application to potatoes 1971 (T):

F1D, F2D, F4D, F6D, F2S, F4S, D2, D4, D2F2D, D2F4D, D4F2D, D4F4D.

D2:D4 = 40: 80 tonnes FYM dug in.

F1 = 126 kg N, 188 kg P2O5, 188 kg K2O.

F2 = 252 kg N, 376 kg P2O5, 376 kg K2O.

F4 = 504 kg N, 752 kg P2O5, 752 kg K2O.

F6 = 756 kg N, 1130 kg P2O5, 1130 kg K2O.

D:S = 'Nitro-Chalk' rotary cultivated in, (0:20:20) dug in:

'Nitro-Chalk' and (0:20:20) raked in shallowly after rotary cultivation.

Sub plots: 2. Nitrogen to winter wheat 1972: 63 kg, 126 kg N as 'Nitro-Chalk'.

Basal applications: Weedkiller: 2,4-D at 0.42 kg with dichlorprop at 1.7 kg in 340 l.

Variety: Cappelle sown at 200 kg.

Cultivations, etc.: Deep-tine cultivated: 5 Oct, 1971. Seed sown: 28 Oct. N applied: 20 Apr, 1972. Weedkiller applied: 4 May. Harvested by hand: 24 Aug.

Standard errors per plot.

Grain, tonnes/hectare: Whole plot: 0.489 or 11.2% (22 d.f.)

Sub plot: 0.416 or 9.6% (24 d.f.)

72/W/CS/77

TABLE OF MEANS

GRAIN: TONNES/HECTARE

T

	F1D	F2D	F4D	F6D	F2S	F4S	D2	D4	D2F2D	D2F4D	D4F2D	D4F4D	Mean
N: KG/HA													
63	3.50	3.86	3.54	3.90	3.67	4.27	4.26	4.58	4.48	3.93	3.83	4.72	4.05
126	4.62	4.98	4.90	4.15	4.23	4.48	4.63	4.78	4.91	4.27	4.65	5.37	4.66
Mean	4.06	4.42	4.22	4.02	3.95	4.37	4.44	4.68	4.69	4.10	4.24	5.05	4.35

227

STANDARD ERRORS OF DIFFERENCES

T	N	T	N
0.399	0.098	0.466	0.340

Unless same levels of T

Mean D.M.  $\phi$ : 82.7

72/W/CS/77

STRAW FRESH: TONNES/HECTARE

T

	F1D	F2D	F4D	F6D	F2S	F4S	D2	D4	D2F2D	D2F4D	D4F2D	F4F4D	Mean
N: KG/HA													
63	7.05	8.45	9.89	7.72	7.58	9.16	9.14	11.39	10.42	8.96	7.72	11.05	9.04
126	7.85	9.39	9.78	7.32	7.39	7.86	8.56	10.47	10.00	8.21	8.17	11.64	8.89
Mean	7.45	8.92	9.83	7.52	7.49	8.51	8.85	10.93	10.21	8.59	7.95	11.35	8.97