

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/W/CS/78 Nematodes and Verticillium - Potatoes

### Rothamsted Research

Rothamsted Research (1973) *72/W/CS/78 Nematodes and Verticillium - Potatoes ; Yields Of The Field Experiments 1972*, pp 229 - 230 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/W/CS/78

NEMATODES AND VERTICILLIUM

Object: To study the residual effects of methyl bromide, aldicarb and benomyl (applied to potatoes in 1971) on *Heterodera rostochiensis* and *Verticillium* on a second potato crop in 1972 - Woburn Broadmead I.

Sponsors: D.C.M. Corbett, G.A. Hide.

For first year see 71/W/CS/78(t).

The second year, potatoes.

Design: 4 blocks of 6 plots.

Whole plot dimensions: 2.84 x 12.8. Area harvested: 0.00147.

Treatments:

Residues of chemicals applied in 1971 (C): None (O), aldicarb at 6.7 kg (A), benomyl at 22.4 kg (B), aldicarb at 6.7 kg plus benomyl at 22.4 kg (A+B), dazomet at 336 kg (D), methyl bromide at 975 kg (M).

Basal applications: 1500 kg (13:13:20), 100 kg MgO as Epsom salts. Weedkiller: Chlorbromuron at 1.7 kg plus paraquat at 0.4 kg ion in 370 l. Fungicide: Mancozeb at 1.3 kg in 430 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 430 l.

Seed: Pentland Dell.

Cultivations, etc.: Deep-tine cultivated: 16 Sept, 1971. NPK applied: 24 Mar, 1972. Mg applied: 17 Apr. Rotary cultivated twice, potatoes planted: 20 Apr. Weedkiller applied: 8 May. Fungicide applied: 6 July. Fungicide with insecticide applied: 27 July. Haulm mechanically destroyed: 24 Aug. Lifted: 12 Sept.

NOTE: Soil samples were taken after harvest for counts of numbers of cysts, eggs and larvae of *Heterodera rostochiensis* and propagules of *Verticillium*.

Standard error per plot.

Total tuber, tonnes/hectare: 3.53 or 21.6 % (15 d.f.)

72/W/CS/78

TABLES OF MEANS

C

O	A	B	A+B	D	M	Mean
TOTAL TUBERS: TONNES/HECTARE						
4.7	20.9	20.9	25.5	9.9	16.1	16.3

STANDARD ERROR OF DIFFERENCES

C 2.49

% WARE: 4.44 CM (1.75 INCH) REDDLE

10.2	63.7	70.2	72.1	26.5	49.5	48.7
------	------	------	------	------	------	------