

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/CS/86 Weedkiller and Aqueous N - Old Grass

Rothamsted Research

Rothamsted Research (1973) *72/R/CS/86 Weedkiller and Aqueous N - Old Grass* ; Yields Of The Field Experiments 1972, pp 235 - 238 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/CS/86

WEEDKILLER AND AQUEOUS N

Object: To study the effects of a range of rates of solid or liquid nitrogen in combination with a range of rates of hormone weedkiller, on weed control and yield of old grass - Ver.

Sponsors: S.C.R. Freeman, A. Penny.

The first year, old grass.

Design: 3 randomised blocks of 28 plots.

Whole plot dimensions: 2.74 x 1.37. Area harvested: 0.00022.

Treatments: (applied per cut): All combinations of:-

1. Weedkiller (dichlorprop/MCPA) (H): None (0), 1.4 (1), 2.8 (2), 4.2 (3) kg total a.e. (using fan jet size 00).
2. Forms of nitrogen: Solid, as 'Nitro-Chalk' 21% N applied immediately after the weedkiller, liquid, as urea/ammonium nitrate (26% N), mixed with the weedkiller (using fan jet size 00).
3. Nitrogen: 37.7, 75.3, 113.0 kg N, together with 4 extra treatments (all liquid N₃):- H0, H1, H2, H3, where fan jet size 1 was used.

NOTE: The weedkiller was applied in 337 l where solid fertiliser was used. The liquid fertiliser (with or without weedkiller) was applied as a spray at 112, 225 and 337 l for rates 1, 2 and 3 respectively.

Basal applications: 630 kg 0:14:28 broadcast.

Cultivations, etc.: Grass lightly trimmed, basal PK broadcast: 10 Mar, 1972. All treatments applied: 13 Apr. Cut: 8 June. All treatments re-applied: 28 June. Cut: 28 Sept.

- NOTES: (1) Scores were made of foliar scorch by treatments on 18 Apr and 6 July.
(2) A score of the presence of weeds was made on 31 May.
(3) The % N in herbage was measured.
(4) The yield of weeds at the second cut was measured.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut:	0.575 or 8.3% (54 d.f.)
2nd cut:	0.451 or 17.0% (54 d.f.)
Total of 2 cuts:	0.873 or 9.1% (54 d.f.)

72/R/CS/86

TABLES OF MEANS

DRY MATTER: TONNES/HECTARE

1ST CUT

	FORM		N:KG/HA			Mean
	Solid	Liquid	37.7	75.3	113.0	
H: KG/HA						
0	7.15	6.68	6.38	7.00	7.36	6.92
1.4	7.32	6.49	6.73	7.05	6.94	6.91
2.8	7.05	6.56	5.94	7.01	7.47	6.81
4.2	6.93	6.66	6.40	6.76	7.22	6.80
		Form				
		Solid	6.53	7.35	7.46	7.11
		Liquid	6.19	6.57	7.04	6.60
Mean			6.36	6.96	7.25	6.86

EXTRA (LIQUID N3)

H: KG/HA					Mean
0	1.4	2.8	4.2		
7.92	7.55	7.36	7.19	7.51	

STANDARD ERRORS OF DIFFERENCES

H	FORM	N	H FORM	H N	FORM N	EXTRA
0.192	0.135	0.166	0.324	0.376	0.235	0.469

Grand Mean: 6.95
Mean D.M.%: 23.3

72/R/CS/86

DRY MATTER: TONNES/HECTARE

2ND CUT

H: KG/HA	FORM		N:KG/HA			Mean
	Solid	Liquid	37.7	75.3	113.0	
0	2.89	2.54	2.14	2.65	3.35	2.72
1.4	3.16	2.27	2.27	2.73	3.15	2.72
2.8	2.77	2.52	2.29	2.58	3.08	2.65
4.2	2.55	2.25	2.00	2.37	2.83	2.40
		Form				
		Solid	2.43	2.68	3.42	2.85
		Liquid	1.91	2.48	2.79	2.39
Mean			2.17	2.58	3.10	2.62

EXTRA (LIQUID N3)

H: KG/HA					Mean
0	1.4	2.8	4.2		
3.53	2.68	2.92	2.36		2.88

STANDARD ERRORS OF DIFFERENCES

H	FORM	N	H FORM	H N	FORM N	EXTRA
0.150	0.106	0.130	0.254	0.296	0.184	0.369

Grand Mean: 2.66
Mean D.M. %: 31.0

72/R/CS/86

DRY MATTER: TONNES/HECTARE

TOTAL OF 2 CUTS

H: KG/HA	FORM		N:KG/HA			Mean
	Solid	Liquid	37.7	75.3	113.0	
0	10.05	9.22	8.53	9.66	10.71	9.63
1.4	10.49	8.76	8.99	9.78	10.09	9.62
2.8	9.82	9.09	8.22	9.59	10.55	9.45
4.2	9.49	8.91	8.39	9.14	10.06	9.20
		Form				
		Solid	8.97	10.04	10.88	9.96
		Liquid	8.10	9.05	9.82	8.99
Mean			8.54	9.54	10.35	9.48

EXTRA(LIQUID N3)

H: KG/HA					Mean
0	1.4	2.8	4.2		
11.46	10.24	10.28	9.55		10.38

STANDARD ERRORS OF DIFFERENCES

H	FORM	N	H FORM	H N	FORM N	EXTRA
0.291	0.206	0.252	0.492	0.572	0.356	0.713

Grand Mean: 9.61

Mean D.M.%: 27.1