

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/S/CS/1 Row Spacing and Seed Rates After Intensive Wheat - W. Wheat

#### Rothamsted Research

Rothamsted Research (1973) *72/S/CS/1 Row Spacing and Seed Rates After Intensive Wheat - W. Wheat* ; Yields Of The Field Experiments 1972, pp 267 - 269 - DOI:

<https://doi.org/10.23637/ERADOC-1-62>

72/S/CS/1

ROW SPACING AND SEED RATES AFTER INTENSIVE WHEAT

Object: To study the effects of seed rate and row spacing on the yields of winter wheat. The effects of different periods of pre-cropping with winter wheat are also studied - Saxmundham Oldershaw's and Garner's plots.

Sponsors: F.V. Widdowson, A.E. Johnston.

The seventh year on this site, second year of row spacing and seed rate treatments, winter wheat.

For previous years see 66/C/30(t), 67/C/23(t), 68/C/39, 69-70/S/CS/1 and 71/S/CS/1(t).

Design: A single replicate of 4 x 2 x 4 in 4 blocks of 4 plots each split lengthways into 2, plus one additional plot (not split into 2) per block. Additionally, all the plots are split breadthways into 3.

Whole plot dimensions: 5.49 x 40.2. Sub plot area harvested: 0.00312.

Treatments: Cumulative on 1971 treatments.

Additional plots: Two immediately previous continuous winter wheat crops (Year 2), row spacing 189 mm (8.25 inches) and seed rate 212 kg (S), and N to sub-plots as shown below, and all combinations of:-

Whole plots:

1. Cropping history: 3, 4, 5 and 6 years of immediately previous winter wheat crops 1966 - 1971 (3, 4, 5, 6).

Half plots:

2. Row spacing: 152 mm (6 inches), 304 mm (12 inches).
3. Seed rates: 70.6, 141, 212, 283 kg.

One third plots:

4. N rates: 50, 100, 150 kg N as 'Nitro-Chalk'.

Basal applications: 1260 kg (0:20:20) applied to stubble in autumn before ploughing. 310 kg (20:10:10) broadcast at drilling.

Weedkiller: Terbutryne and related triazines ('Prebane' at 4.5 kg).

Seed: Cappelle.

72/S/CS/1

Cultivations, etc.: Basal PK applied and ploughed in: 2 Oct, 1971.  
Seed drilled and basal NPK applied: 26 Oct. Weedkiller applied:  
27 Oct. Test N applied: 21 Apr, 1972. Combine harvested:  
29 Aug.

Addendum to 71/S/CS/1: Basal applications should include 310 kg  
(20:10:10) broadcast at drilling.

NOTE: Green crop samples for estimates of total dry matter and leaf  
areas were taken.

72/S/CS/1

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	Year				Mean
	3	4	5	6	
<b>SPACING: MM</b>					
152	5.82	5.80	5.92	5.90	5.86
304	5.93	5.88	5.86	5.85	5.88
<b>SEED RATE: KG</b>					
70.6	5.95	5.71	5.85	5.82	5.83
141	5.96	5.96	5.98	5.93	5.96
212	5.82	5.98	5.96	5.98	5.93
283	5.77	5.71	5.76	5.77	5.75
<b>N: KG/HA</b>					
50	5.38	5.35	5.13	5.17	5.26
100	6.07	6.16	6.16	6.25	6.16
150	6.17	6.01	6.37	6.21	6.19
Mean	5.87	5.84	5.89	5.87	5.87

YEAR 2

N: KG/HA

50	100	150	Mean
4.31	4.83	5.87	5.34

Grand mean: 5.76  
Mean D.M. %: 84.8