

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/R/WW/1 and 72/W/WW/1 Winter Wheat Varieties and N

### Rothamsted Research

Rothamsted Research (1973) *72/R/WW/1 and 72/W/WW/1 Winter Wheat Varieties and N* ; Yields Of The Field Experiments 1972, pp 270 - 273 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/1 and 72/W/WW/1

WINTER WHEAT

VARIETIES AND N

**Object:** To study the yield of newer varieties of wheat grown at a range of nitrogen levels, on land in rotation or after several cereals. Nitrogen applied at flowering is also tested, and bread-making quality is determined. Rothamsted (R) Whittlocks (pathogen free) and Claycroft (pathogen infected) and Woburn (W) Great Hill II (pathogen free).

**Sponsors:** J.R. Moffatt, R. Moffitt.

**Design:** 3 randomised blocks of 8 plots, split into 4.

**Whole plot dimensions:** 4.27 x 27.1. **Sub plot area harvested:** 0.00173.

**Treatments:** All combinations of:-

**Whole plots:** 1. Varieties: Champlain (CH), Cama (CM), Cappelle (CP), Joss Cambier (JC), Maris Huntsman (MH), Maris Nimrod (MN), Maris Widgeon (MW), Tommy (TO).

**Sub plots:** 2. Levels of nitrogen (in kg N): 63, 126, 189 in spring, and 126 in spring plus 63 at flowering. All N as 'Nitro-Chalk'.

**Basal applications:**

**Manures:** Claycroft (R) and Whittlocks (R): 300 kg (0:20:20).

Great Hill II (W): 290 kg (0:20:20).

**Weedkillers:** Paraquat at 0.56 kg ion in 220 l to Claycroft (R) only. 2,4-D at 0.7 kg plus dichlorprop at 2.8 kg in 220 l (280 l on Great Hill II (W)) to Claycroft (R) and Great Hill II (W). MCPA, mecoprop and dicamba ('Banlene Plus' at 5.6 l in 220 l) to Whittlocks (R).

**Seed:** Varieties sown at 200 kg on Claycroft (R) and Whittlocks (R) and at 180 kg on Great Hill II (W).

**Cultivations, etc.:**

Claycroft (R): Paraquat applied: 28 Sept, 1971. Deep-tine cultivated on 3 occasions: 12, 22 and 25 Oct. Seed combine drilled: 27 Oct. N applied: 5 Apr, 1972. Weedkiller applied: 26 Apr. Late N applied: 15 June. Combine harvested: 30 Aug. Previous crops: Winter wheat 1970 and 1971.

72/R/WW/1 and 72/W/WW/1

Whittlocks (R): Deep-tine cultivated on 2 occasions: 25 and 26 Oct, 1971. Seed combine drilled: 28 Oct. N applied: 5 Apr, 1972. Weedkiller applied: 27 Apr. Late N applied: 15 June. Combine harvested: 30 Aug. Previous crops: Fallow 1970, potatoes 1971.

Great Hill II (W): Deep-tine cultivated on 2 occasions: 8 and 9 Oct, 1971. Seed combine drilled: 22 Oct. N applied: 17 Apr, 1972. Weedkiller applied: 27 Apr. Late N applied: 19 June. Combine harvested: 30 Aug. Previous crops: Fallow 1970, potatoes 1971.

Standard errors per plot. Grain, tonnes/hectare:

Whittlocks (R):	Whole plot: 0.398 or 6.0% (14 d.f.)
	Sub plot: 0.740 or 11.1% (48 d.f.)
Claycroft (R):	Whole plot: 0.449 or 7.3% (14 d.f.)
	Sub plot: 0.433 or 7.0% (44 d.f.)
Great Hill II (W):	Whole plot: 0.449 or 10.9% (14 d.f.)
	Sub plot: 0.444 or 10.8% (48 d.f.)

NOTE: On Claycroft (R) four plots of one block were damaged by cattle and yields were not taken. Estimated values were used in the analysis. The treatment combinations affected were:-

Variety	N
TO	126
CH	63
CM	63
MN	63



72/R/WW/1 and 72/W/WW/1

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	VARIETY								Mean
	CH	CM	CP	JC	MH	MN	MW	TO	
WHITTLOCKS (R) PATHOGEN FREE									
N: KG/HA									
63	7.27	6.24	6.21	5.81	6.74	7.44	5.92	6.35	6.50
126	6.12	6.82	6.19	4.67	8.40	8.41	6.40	7.23	6.78
189	6.31	4.80	6.08	4.14	8.22	7.49	6.89	5.84	6.22
126+63	7.60	6.19	7.63	4.71	8.37	8.33	6.98	6.88	7.09
Mean	6.82	6.01	6.53	4.83	7.93	7.91	6.55	6.58	6.65

STANDARD ERRORS OF DIFFERENCES

VARIETY	NITROGEN	VARIETY NITROGEN
0.325	0.214	0.616
Unless same levels of VARIETY		0.604

Mean D.M. %: 84.1

	CLAYCROFT (R) PATHOGEN INFECTED								Mean
	CH	CM	CP	JC	MH	MN	MW	TO	
N: KG/HA									
63	5.62	5.24	5.28	5.35	6.06	6.80	5.55	5.71	5.70
126	6.56	5.68	5.91	5.58	7.11	7.02	6.20	6.60	6.33
189	6.13	5.71	5.54	4.80	7.32	7.52	6.36	5.89	6.16
126+63	7.09	5.59	5.86	5.49	7.81	7.70	6.03	6.50	6.51
Mean	6.35	5.56	5.65	5.30	7.07	7.26	6.03	6.17	6.17

STANDARD ERRORS OF DIFFERENCES

VARIETY	NITROGEN	VARIETY NITROGEN
0.366	0.125	0.477
Unless same levels of VARIETY		0.355

Mean D.M. %: 85.4

272

72/R/WW/1 and 72/W/WW/1

GRAIN: TONNES/HECTARE

GREAT HILL II (W) PATHOGEN FREE

	VARIETY								Mean
	CH	CM	CP	JC	MH	MN	MW	TO	
N: KG/HA									
63	4.16	3.62	4.85	3.66	4.33	4.65	4.06	4.15	4.19
126	4.75	3.10	4.99	3.69	4.55	4.42	4.69	4.02	4.28
189	5.07	2.73	3.84	2.96	3.54	4.61	3.90	3.22	3.73
126+63	5.25	3.38	4.62	3.36	4.62	4.70	4.51	3.58	4.25
Mean	4.81	3.21	4.57	3.42	4.26	4.59	4.29	3.74	4.11

STANDARD ERRORS OF DIFFERENCES

VARIETY	NITROGEN	VARIETY NITROGEN
0.367	0.128	0.483
Unless same levels of VARIETY		0.363

Mean D.M. %: 85.8