

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1972

[Full Table of Content](#)



## 72/R/WW/2 Winter Wheat Weedkiller and Aqueous N

### Rothamsted Research

Rothamsted Research (1973) 72/R/WW/2 Winter Wheat Weedkiller and Aqueous N ; Yields Of The Field Experiments 1972, pp 274 - 276 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/2

WINTER WHEAT

WEEDKILLER AND AQUEOUS N

Object: To study the effects of a range of rates of solid or liquid nitrogen in combination with a range of rates of hormone weedkiller on foliar scorch and yield of winter wheat - Whittlocks.

Sponsors: S.C.R. Freeman, A. Perry.

Design: 3 randomised blocks of 28 plots.

Whole plot dimensions: 2.13 x 2.74. Area harvested: 0.00038.

Treatments: All combinations of:-

1. Weedkiller (dichlorprop/MCPA) (H): None (0), 1.4 (1), 2.8 (2), 4.2 (3) kg total a.e.
2. Forms of nitrogen: Solid, as 'Nitro-Chalk' (21% N) applied immediately after the weedkillers, liquid, as urea/ammonium nitrate (26% N) mixed with the weedkiller (L).
3. Nitrogen: 37.7, 75.3, 113.0 kg N, together with 4 extra treatments: SN2 E HO, SN2 E HL, SN2 E H2, SN2 E H3 (N2 = 75.3 kg N) where 'Nitro-Chalk' was applied early (E) and the HO plots were hand weeded.

Basal applications: 280 kg (0:20:20) combine drilled.

Seed: Joss Cambier dressed dieldrin, sown at 200 kg.

Cultivations, etc.: Deep-tine cultivated twice: 28 Oct, 1971. Seed combine drilled: 29 Oct. N applied to E plots: 17 Apr, 1972. Remaining N treatments and weedkiller applied: 2 May. HO plots hand weeded: 30 June. Cut by sickle: 29 Aug. Previous crops: Fallow 1970, potatoes 1971.

NOTE: Grain and straw samples were taken for determination of dry matter and samples retained for N analyses.

Standard error per plot.

Grain, tonnes/hectare: 0.403 or 7.3% (54 d.f.)

72/R/WW/2

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	H: KG/HA				Mean
	0	1.4	2.8	4.2	
FORM					
S	5.45	5.70	5.44	5.54	5.53
L	5.32	5.70	5.54	5.26	5.46
N: KG/HA					
37.7	5.24	5.26	5.18	5.24	5.23
75.3	5.43	6.11	5.73	5.39	5.67
113.0	5.50	5.72	5.56	5.57	5.59
Mean	5.39	5.70	5.49	5.40	5.49

	N: KG/HA		
	37.7	75.3	113.0
FORM			
S	5.39	5.72	5.49
L	5.07	5.61	5.68

EXTRA (SN2 E )

H: KG/HA	
0	5.28
1.4	5.47
2.8	6.20
4.2	5.84

STANDARD ERROR OF DIFFERENCES

H	FORM	N	H FORM	H N	EXTRA
0.134	0.095	0.116	0.227	0.264	0.329

Grand mean: 5.52

Mean D.M. %: 83.8



72/B/WW/2

STRAW: TONNES/HECTARE

	H: KG/HA				Mean
	0	1.4	2.8	4.2	
FORM					
S	8.32	8.19	8.16	8.06	8.18
L	7.89	7.93	7.80	7.34	7.74
N: KG/HA					
37.7	7.66	7.45	6.83	7.01	7.24
75.3	7.84	8.48	8.17	7.58	8.02
113.0	8.80	8.25	8.93	8.52	8.63
Mean	8.10	8.06	7.98	7.70	7.96

N: KG/HA

	37.7	75.3	113.0
FORM			
S	7.34	8.27	8.92
L	7.13	7.76	8.33

EXTRA (SN2 E)

H:KG/HA	
0	8.68
1.4	8.58
2.8	8.63
4.2	8.60

Grand mean: 8.05

Mean D.M. %: 69.0