

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1972

[Full Table of Content](#)



### 72/R/WW/3 Winter Wheat Weedkiller, Aqueous N and Fungicide

#### Rothamsted Research

Rothamsted Research (1973) *72/R/WW/3 Winter Wheat Weedkiller, Aqueous N and Fungicide ; Yields Of The Field Experiments 1972*, pp 277 - 279 - DOI:

<https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/3

WINTER WHEAT

WEEDKILLER, AQUEOUS N AND FUNGICIDE

Object: To study the effects of combined sprays of liquid nitrogen fertiliser, hormone weedkiller and fungicide on foliar scorch and yield of winter wheat - Whittlocks.

Sponsors: S.C.R. Freeman, A. Penny.

Design: 4 randomised blocks of 21 plots.

Whole plot dimensions: 2.13 x 2.74. Area harvested: 0.00264.

Treatments: All combinations of:-

1. Forms of nitrogen: Solid as 'Nitro-Chalk' (S) (21% N) applied immediately after weedkiller and fungicide, liquid as urea/ammonium nitrate (26% N) mixed with the weedkiller and fungicide where appropriate.

2. Nitrogen: 56.5, 113 kg N.

3. Weedkiller (dichlorprop/MCPA) (H): None, 2.8 kg total a.e.

4. Fungicide (F): None (O), tridemorph at 0.53 kg (T)

together with 5 extra treatments:-

SN1E H TO, SN2E H TO, SN1E H TE, SN2E H TE, SN2 H TL,

(N1 = 56.5, N2 = 113.0 kg N, H = H at 2.8 kg)

where 'Nitro-Chalk' and tridemorph were applied early (E), and tridemorph was applied late (L).

Basal applications: 280 kg (0:20:20) combine drilled.

Seed: Joss Cambier dressed dieldrin, sown at 200 kg.

Cultivations, etc.: Deep-tine cultivated twice: 28 Oct, 1971. Seed combine drilled: 29 Oct. N and tridemorph applied to E plots: 17 Apr, 1972. Remaining N and tridemorph treatments and weedkiller applied: 11 May. Cut by sickle: 30 Aug. Previous crops: Fallow 1970, potatoes 1971.

NOTE: Estimates were made of infection by mildew (*Erysiphe graminis*). Grain samples were taken for determinations of dry matter and % N.

Standard error per plot.

Grain, tonnes/hectare: 0.375 or 6.9% (60 d.f.)

72/R/WW/3

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	N: KG/HA		H: KG/HA		F: KG/HA		Mean
	56.5	113.0	0	2.8	0	0.53	
FORM							
Solid	5.56	5.30	5.41	5.45	5.39	5.47	5.43
Liquid	5.30	5.22	5.19	5.34	5.42	5.11	5.26
	N: KG/HA		H: KG/HA		F: KG/HA		
	56.5	113.0	5.36	5.51	5.52	5.35	5.43
			5.24	5.27	5.29	5.23	5.26
			H: KG/HA		F: KG/HA		
			0	2.8	5.32	5.28	5.30
					5.49	5.30	5.39
Mean					5.40	5.29	5.35

EXTRA

SN1E H TO SN2 H TO SN1E H TE SN2 H TE SN2 H TL

5.79 5.54 5.86 5.30 5.45

STANDARD ERRORS OF DIFFERENCES

ALL MARGINAL MEANS ALL 2-FACTOR TABLES EXTRA

0.094 0.133 0.265

Grand mean: 5.40

Mean D.M. %: 84.9

72/R/WW/3

STRAW (FRESH): TONNES/HECTARE

	N: KG/HA		H: KG/HA		F: KG/HA		Mean
	56.5	113.0	0	2.8	00	0.53	
FORM							
Solid	9.81	9.40	9.87	9.34	9.54	9.67	9.60
Liquid	9.27	9.27	9.63	8.91	9.49	9.06	9.27
	N: KG/HA						
	56.5		9.94	9.15	9.50	9.58	9.54
	113.0		9.56	9.11	9.53	9.14	9.34
			H: KG/HA				
			0		9.82	9.69	9.75
			2.8		9.21	9.04	9.13
Mean					9.51	9.36	9.44

EXTRA

SN1E H TO SN2 H TO SN1E H TE SN2E H TE SN2 H TL

10.89 10.61 11.08 11.17 9.29

Grand mean: 9.72