Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1972



Full Table of Content

72/R/WW/4 Winter Wheat Ewsm Spread by Infected Vectors

Rothamsted Research

Rothamsted Research (1973) 72/R/WW/4 Winter Wheat Ewsm Spread by Infected Vectors; Yields Of The Field Experiments 1972, pp 280 - 281 - DOI: https://doi.org/10.23637/ERADOC-1-62

72/R/WW/4

WINTER WHEAT

EWSM SPREAD BY INFECTED VECTORS

Object: To study the effects of introducing vectors (Javesella pellucida), carrying european wheat striate mosaic, at different times, on spread of disease and yield of wheat - Long Hoos IV.

Sponsor: R.T. Plumb.

Design: 2 randomised blocks of 5 plots.

Whole plot dimensions: 6.40 x 12.2. Area harvested: 0.00347.

Treatments: Caging of plant-hopper vectors (Javesella pellucida) carrying EWSM on central area (0.61 x 1.22) of plot for 2 weeks (DATE):-

No vectors

Vectors caged on 23 December, 1971

On 20 January, 1972

On 5 April

On 8 May

(M)

The central area (0.61 x 1.22) was cut out before plot yields were taken.

Basal applications: 250 kg (10:24:24) combine drilled. 105 kg N as 'Nitro-Chalk'. Weedkiller: Dicamba, mecoprop and MCPA ('Tetralex Plus' at 7 l in 220 l).

Seed: Cappelle sown at 200 kg.

Cultivations, etc.: Ploughed: 6 Oct, 1971. Seed combine drilled: 23 Oct. N applied: 24 Apr, 1972. Weedkiller applied: 26 Apr. Combine harvested: 31 Aug. Previous crops: Potatoes 1970, barley 1971.

NOTE: Scores for EWSM were made and plants were sampled throughout the season.

Standard error per plot.

Grain, tonnes/hectare: 0.149 or 2.0% (4 d.f.)

280

72/R/WW/4

TABLE OF MEANS

GRAIN: TONNES/HECTARE

DATE

0	D	J	A	М	Mean
7.36	7-35	7.38	7.11	7.31	7.30

STANDARD ERROR OF DIFFERENCES
DATE 0.149

Mean D.M. %: 85.0