

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/WW/6 and 72/BB/WW/6 Winter Wheat Growth and Yield on Contrasted Sites

Rothamsted Research

Rothamsted Research (1973) *72/R/WW/6 and 72/BB/WW/6 Winter Wheat Growth and Yield on Contrasted Sites* ; Yields Of The Field Experiments 1972, pp 282 - 285 - DOI:

<https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/6 and 72/BB/WW/6

WINTER WHEAT

GROWTH AND YIELD ON CONTRASTED SITES

Object: To try to account for yields and differences between yields of wheat on sites at Rothamsted and Broom's Barn by studying crop growth rates, nutrient uptake, water use etc., at a wide range of nitrogen levels, with and without irrigation. Also to study the interaction between site differences and crops (see also 72/R&BB/B/1) - Rothamsted (R), Long Hoos I and II and Broom's Barn (BB), Hackthorn Field.

Sponsors: P.J. Welbank, F.V. Widdowson.

Design: 3 randomised blocks of 2 plots, split into 6.

Whole plot dimensions:-

Long Hoos I and II (R): 15.2 x 47.8. Sub plot area harvested: 0.00434.
Hackthorn Field (BB): 15.2 x 45.7. Sub plot area harvested: 0.00413.

Treatments: All combinations of:-

Whole plots: 1. Irrigation (I): None (O), full irrigation (C).

Sub plots: 2. Nitrogen: 31, 63, 94, 125, 157, 188 kg N as 'Nitro-Chalk'.

Total irrigation was 86.4 mm, applied on 4 occasions (R) and 126 mm, on 5 occasions (BB).

Basal applications (both fields):

1260 kg (0:20:20) plus 900 kg Epsom salts in seedbed. 250 kg (10:24:24) combine drilled. Weedkiller: MCPA, mecoprop and dicamba ('Banlene Plus' at 5.6 l in 220 l). Fungicide: Tridemorph at 0.53 kg in 280 l.

Seed: Cappelle, dressed with dieldrin, sown at 200 kg.

Cultivations, etc.:

Long Hoos I and II (R): Basal PK and Epsom salts applied:

24 Sept, 1971. Ploughed: 5 Oct. Seed combine drilled:

25 Oct. N applied: 19 Apr, 1972. Weedkiller applied:

26 Apr. Fungicide applied: 12 May. Irrigated: 2, 13 and

29 June and 14 July. Combine harvested: 31 Aug. Previous

crops: Spring wheat 1970, spring beans 1971.

72/R/WW/6 and 72/BB/WW/6

Hackthorn Field (BB): Deep-tine cultivated: 23 Sept, 1971.
Epsom salts applied: 12 Oct. Basal PK applied: 15 Oct.
Seed combine drilled: 25 Oct. Weedkiller applied: 17 Apr.
N applied: 19 Apr, 1972. Fungicide applied: 16 May.
Irrigated: 2, 13, 29 and 30 June and 18 July. Combine
harvested: 24 Aug. Previous crops: Sugar beet 1970,
beans 1971.

Standard errors per sub plot. Grain, tonnes/hectare:

Long Hoos I and II (R): 0.272 or 3.6% (20 d.f.)

Hackthorn Field (BB): 0.432 or 7.0% (20 d.f.)

72/R/WW/6 and 72/BB/WW/6

TABLES OF MEANS

LONG HDOS I & II (R)

N: KG/HA

	31	63	97	125	157	188	Mean
I	GRAIN: TONNES/HECTARE						
D	8.04	8.24	7.86	7.68	7.50	6.25	7.59
C	7.84	7.99	8.02	7.59	7.48	6.72	7.61
Mean	7.94	8.11	7.94	7.64	7.49	6.48	7.60

STANDARD ERRORS OF DIFFERENCES

N	I* N
0.157	0.222

* Within the same level of I

I	STRAW: TONNES/HECTARE						
D	6.90	7.09	7.53	7.92	8.62	8.35	7.74
C	6.37	7.77	7.82	8.56	8.72	8.57	7.97
Mean	6.63	7.43	7.68	8.24	8.67	8.46	7.85

Mean D.M. %: Grain: 80.9
Straw: 80.9

72/R/WW/6 and 72/BB/WW/6

HACKTHORN FIELD (BB)

N: KG/HA

	31	63	94	125	157	188	Mean
I	GRAIN: TONNES/HECTARE						
O	5.46	6.05	6.37	5.89	6.06	5.85	5.94
C	6.09	6.67	6.71	6.64	6.48	6.14	6.45
Mean	5.77	6.36	6.54	6.26	6.27	5.99	6.20

STANDARD ERRORS OF DIFFERENCES

N	I* N
0.250	0.353

* Within the same level of I

I	STRAW: TONNES/HECTARE						
O	6.53	7.20	7.63	7.23	7.89	7.56	7.34
C	6.18	7.16	7.42	7.66	7.88	7.64	7.32
Mean	6.35	7.18	7.52	7.45	7.88	7.60	7.33

Mean D.M. %: Grain: 85.2
Straw: 60.0