

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/WW/7 Winter Wheat Methods of Applying Fungicides

Rothamsted Research

Rothamsted Research (1973) 72/R/WW/7 *Winter Wheat Methods of Applying Fungicides* ; Yields Of The Field Experiments 1972, pp 286 - 287 - DOI: <https://doi.org/10.23637/ERADOC-1-62>

72/R/WW/7

WINTER WHEAT

METHODS OF APPLYING FUNGICIDES

Object: To study the effects of fungicides, applied to seed or soil, on yields and pathogens of wheat - West Barnfield II.

Sponsors: J. Jenkyn, R.D. Prew.

Design: 3 randomised blocks of 5 plots.

Whole plot dimensions: 2.13 x 13.4. Area harvested: 0.00195.

Treatments: Fungicide (F)

None

(O)

Seed dressings: Organo-mercury fungicide at 425 g of
the commercial formulation
NF48 at 2.28 kg a.i.

(M)

(N)

Soil drench after sowing:

Sandoz fungicide at 11.1 l in 5600 l
and at 16.6 l in 5600 l

(S1)

(S2)

Basal applications: 250 kg (10:24:24) combine drilled, and 600 kg
'Nitro-Chalk' in spring. Weedkillers: Paraquat at 0.42 kg ion
in 220 l. MCPA, mecoprop and dicamba ('Banlene Plus' at 5.6 l
in 220 l).

Seed: Cama (undressed apart from treatments M and N) sown at 190 kg.

Cultivations, etc.: Paraquat applied: 24 Sept, 1971. Ploughed: 9 Oct.
Seed combine drilled: 28 Oct. Soil drench applied: 30 Oct.
N applied: 24 Apr, 1972. 'Banlene Plus' applied: 28 Apr. Combine
harvested: 26 Aug. Previous crops: Winter wheat 1970, barley 1971.

NOTE: Seedling emergence counts were made in January, and samples taken
for root diseases in April and July. Mildew (*Erysiphe graminis*)
was assessed in May, June and July, and yellow rust (*Puccinia
striiformis*) in June and July.

Standard error per plot.

Grain, tonnes/hectare: 0.426 or 11.1% (8 d.f.)

72/R/WW/7

TABLE OF MEANS

GRAIN: TONNES/HECTARE

F					
O	M	N	SL	S2	Mean
4.25	3.80	3.64	3.75	3.83	3.85

STANDARD ERROR OF DIFFERENCES

F 0.348

Mean D.M. %: 85.8